

JOURNAL PRE-PROOF



PARADIGM SHIFT IN UNDERSTANDING THE YARLUNG-TSANGPO OPHIOLITES: INSIGHTS FROM COMPARISONS WITH ULTRASLOW-SPREADING OCEAN RIDGES

Tong Liu, Chuan-Zhou Liu, Chang Zhang, Zhen Zhang, Zhen-Yu Zhang, Wei-Qi Zhang, Yang Xu, Wen-Bin Ji, Fu-Yuan Wu

To appear in OFIOLITI

DOI: <https://doi.org/10.4454/ofioliti.v51i2.581>

Received date: 1 October 2025
Accepted date: 12 January 2026
First published online 17 March 2026

PARADIGM SHIFT IN UNDERSTANDING THE YARLUNG-TSANGPO OPHIOLITES: INSIGHTS FROM COMPARISONS WITH ULTRASLOW-SPREADING OCEAN RIDGES

Tong Liu^{1}, Chuan-Zhou Liu^{1,2,3*}, Chang Zhang¹, Zhen Zhang², Zhen-Yu Zhang⁴, Wei-Qi Zhang⁵, Yang Xu², Wen-Bin Ji⁶, and Fu-Yuan Wu^{1,3}*

¹ State Key Laboratory of Lithospheric and Environmental Coevolution, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China

² Laoshan Laboratory, Qingdao 266237, China

³ College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

⁴ Key Laboratory of Marine Mineral Resources, Ministry of Natural Resources, Guangzhou Marine Geological Survey, China Geological Survey, Guangzhou 511458, China

⁵ Key Laboratory of Submarine Geosciences, Second Institute of Oceanography, Ministry of Natural Resources, Hangzhou 310012, China

⁶ Department of Geology, Northwest University, Xi'an 710069, China

*Corresponding authors: liutong@mail.iggcas.ac.cn (T. Liu); chzliu@mail.iggcas.ac.cn (C.Z. Liu)

Keywords: Yarlung-Tsangpo ophiolites; Tibetan Plateau; Ultraslow-spreading ridge; Ophiolite conundrum; Asthenospheric heterogeneity; Subduction jump and re-initiation

ABSTRACT

The Yarlung-Tsangpo ophiolites (YTO) are crucial tectonic windows that preserve coherent remnants of the Neo-Tethyan oceanic lithosphere in the Tibetan Plateau. They are particularly characterized by geological features that are different from the Penrose model, but resemble oceanic lithospheres generated at slow- to ultraslow-spreading ridges (i.e., the Chapman model). However, geochemical studies on the YTO have widely accepted an origin of suprasubduction zones (SSZ). The seemingly paradoxes between geological and geochemical evidence exemplify the “ophiolite conundrum” that have also been commonly observed in global ophiolites, which thus should be reconciled by any model. The YTO, especially the Xigaze ophiolite in its central segment, exhibits laterally highly variable magmatic crustal sequences, indicating magma focusing during its formation. Notably, the lower crust is either absent or formed through episodic and intermittent magma supplies, probably representing the low end of the magma budget among ocean ridges over time. Decoupled with the thin crust, the mantle peridotites show significant compositional variations (e.g., bulk Al₂O₃ of 0.04–2.9 wt.% and spinel Cr/(Cr + Al) of 0.12–0.74) and reveal heterogeneous osmium isotope compositions (187Os/188Os of 0.11295–0.14661). Such features indicate that they represent mantle residues of a heterogeneous asthenosphere containing recycled ancient mantle domains of variable origins. These findings, combined with the widespread occurrence of detachment faults, support an origin of ultraslow-spreading ridges for the YTO. In this scenario, the ophiolite conundrum of the YTO can be elucidated through alternative models: 1) re-melting of ancient mantle domains within the heterogeneous asthenosphere, and 2) subduction jump and re-initiation along detachment faults, a process superimposed arc-like magmatism on pre-existing mid-ocean-ridge-basalt-like oceanic crusts. Therefore, we suggest that ancient and modern oceanic lithospheres generated at ultraslow-spreading ridges provide resolutions to the “ophiolite

conundrum” and also call upon a paradigm shift in our understanding of global ophiolites from the Penrose model to the Chapman model.

INTRODUCTION

Ophiolites are a suite of mafic–ultramafic rocks that are widely identified in global orogenic belts and suture zones (Anonymous, 1972). They are widely accepted as ancient oceanic lithospheres that have formed in variable tectonic settings, such as mid-ocean ridge (MOR), suprasubduction zone (SSZ), and ocean–continent transition (OCT) (e.g., Dilek and Furnes, 2011). Ophiolites distributed along the Yarlung-Tsangpo suture (YTS) in the southern Tibetan Plateau are the Early Cretaceous remnants of the Neo-Tethys Ocean, an ancient ocean that existed between the Gondwana and Eurasian continents during the Mesozoic (Hébert et al., 2012; Liu et al., 2022b; Wu et al., 2014; Zhu et al., 2013). These ophiolites serve as crucial geological records for deciphering the seafloor structures and evolutionary histories of the Neo-Tethys Ocean, providing valuable insights into regional plate configurations. The Yarlung-Tsangpo ophiolites (YTO) have been the focus of multidisciplinary researches for more than sixty years since the pioneering work of Gansser (1964) and Chang and Zheng (1973). However, the origin of the YTO, in particular its tectonic setting of formation (i.e., MOR or SSZ) remains hotly debated, with diverse models proposed.

The tectonic setting of the YTO was first discussed during a joint China–France fieldwork in the early 1980s (Girardeau et al., 1985a, b; Nicolas et al., 1981; Wang et al., 1987). Systematic field observations in that joint fieldwork revealed a series of “peculiar” geological features in the central part of the YTO, which can be summarized as: 1) a thick (up to > 10 km) mantle unit accompanied by a thin (< 2–3 km) crust, and 2) the absence of a sheeted dike complex replaced instead by isolated and sheeted diabase sills (Girardeau et al., 1985b; Nicolas et al., 1981). Together with the rare occurrence of lower crustal unit, these “peculiar” geological features, which are quite different from the classical “Penrose model”, were considered as key evidence supporting a slow-spreading ridge origin for these ophiolites (Nicolas et al., 1981). In contrast, geochemical studies over the last two decades have identified arc-like signatures in both the basaltic rocks and mantle peridotites of the YTO, leading to an argument that the YTO formed in a SSZ setting, either in a forearc or back-arc context (e.g., Dai et al., 2013, 2011; Dubois-Côté et al., 2005; Hébert et al., 2012; Malpas et al., 2003). Similar hypotheses have further been proposed based on geochronological and geochemical studies of metamorphic soles underlying the YTO, demonstrating a rapid tectonic transition from mid-ocean ridges to back-arc basins (Bédard et al., 2009; Guilmette et al., 2009, 2012). As a result,

the understanding of the tectonic setting of the YTO has transformed significantly from a slow-spreading ocean ridge to subduction-related settings. The YTO is now regarded more as a geological record of subduction initiation within the Neo-Tethys Ocean, rather than a product of seafloor spreading at mid-ocean ridges.

Regardless of its tectonic setting, one of the major achievements in understanding the YTO is the identification of detachment faults and oceanic core complexes (Li et al., 2016; Liu et al., 2014; Maffione et al., 2015; Wu et al., 2014), which are diagnostic geological features of modern slow- and ultraslow-spreading ocean ridges, i.e., the Chapman model (Escartín and Canales, 2011; Maffione et al., 2013). The presence of detachment faults within the YTO suggests a relatively low magma supply during its formation (Tucholke et al., 2008). This geological observation goes against the high magma supply typically expected in a forearc setting owing to large-degree, multi-stage mantle melting (Ishizuka et al., 2011; Stern et al., 2012). Moreover, a classical forearc crust, which is typically composed from the bottom up of gabbros, forearc basalts, boninites, and subsequent island arc basalts, is notably absent in the YTO (Wu et al., 2014). Therefore, whether a MOR or a SSZ origin is considered, any proposed model must reconcile the evidence of a low magma supply, the development of detachment faults, and the observed arc-like geochemical features.

Here we present an updated review of our findings regarding the origin of the YTO over the past decade since the work by Liu et al. (2014) and Wu et al. (2014). Along with the compilation and re-evaluation of published geochemical data, the geological and geochemical characteristics of the YTO will be compared comprehensively with the new findings of modern slow- and ultraslow-spreading ridges. In this context, the arc-like geochemical signatures observed in the YTO can be explained by alternative models, such as the re-melting of ancient mantle domains in the heterogeneous asthenosphere and a subduction re-initiation superimposed on mid-ocean ridge processes. These models, inspired by modern slow- and ultraslow-spreading ridges, provide new insights into the long-standing controversy on the tectonic setting of the YTO. They also have a potential to resolve the “ophiolite conundrum”, which describes the discrepancy between MOR-type structural and stratigraphic features and SSZ-type geochemical characteristics observed in global ophiolites (Moore et al., 2000, 2021), and call upon a paradigm shift in the understanding of global ophiolites from the Penrose model to the Chapman model.

GEOLOGICAL BACKGROUND

Tectonic framework of the Tibetan Plateau

The Tibetan Plateau has formed through the tectonic amalgamation of several continental and/or accretionary terranes, which, from north to south, include the Kunlun-Qilian, Songpan-Ganzi, North and South Qiangtang, and Lhasa (Fig. 1a; Yin and Harrison, 2000; Zhu et al., 2013). These terranes are separated by a series of east–west trending suture zones of the Tethys Ocean, namely the A’nyemaqen-Kunlun-Muztagh, Jinsha, and Longmuco-Shuanghu sutures of the Paleotethys, the Bangong-Nujiang suture of the Meso-Tethys, and the YTS of the Neo-Tethys (Fig. 1a; Wu et al., 2020; Zhu et al., 2013). The YTS is the youngest and southernmost suture within the Tibetan Plateau, marking the surface boundary between the Lhasa terrane to the north and the Himalaya (Indian Plate) to the south.

Along the YTS, several tectonic units have been well identified in relation to the Neo-Tethyan evolution (Fig. 1b). These units include, from north to south, the Triassic–Eocene Gangdese batholith and Linzizong volcanics (known as the Gangdese continental arc), the Early Cretaceous–Eocene Xigaze forearc basin, the YTO, the Triassic–Eocene accretionary and tectonic mélanges. Additionally, terranes with debated origins, such as the Zhongba terrane and Langjiexue Group, have also been identified in southern Tibet.

Geology of the Yarlung-Tsangpo ophiolites

Ophiolites crop out discontinuously along the YTS and extend for more than 1000 km (Fig. 1b; Dai et al., 2013; Hébert et al., 2012; Liu et al., 2022b; Wu et al., 2014; Xiong et al., 2016; Xu et al., 2020). These ophiolites are geographically divided into the western, central, and eastern segments (Fig. 1b). The western YTO is further grouped into northern (Dajiweng, Baer, Cuobuzha, and Zhalai) and southern (Dongbo, Purang, Zhagabu, Mayum, Danqiong, Xiugugabu, Zhongba, and Saga) sub-belts, which are separated by the Zhongba terrane. These ophiolites are mainly made up of variably serpentized mantle peridotites that are surrounded by chert-matrix mélanges, whereas the magmatic crusts are extremely thin or even missing (Bédard et al., 2009; Bezard et al., 2011; Lian et al., 2016; Liu et al., 2014, 2018a; Zhang et al., 2023a, 2023d, 2025). Ophiolites in the central segment are distributed mainly south of Xigaze City, stretching from Ngamring in the west to Renbu in the east, thus are combined referred to as the Xigaze ophiolite. As the most studied ophiolite in China, the Xigaze ophiolite has been the focus of extensive research, regarding its field geology, structure, geochronology, petrology, geochemistry, and paleomagnetism. The Xigaze ophiolite has a complete and continuous sequence, from the uppermost mantle in the south, through

the lower crust, to upper crustal basaltic rocks in the north (Girardeau et al., 1985a, b; Liu et al., 2021a, b; Nicolas et al., 1981; Wu et al., 2014). The detailed geological features and tectonic significance of the Xigaze ophiolite are provided in the Discussion section. Ophiolites in the eastern segment, mainly located in Zedong and Luobusa, exhibit geological features that differ significantly from those in the western and central segments. The eastern ophiolites typically consist of a mantle sequence of lherzolites and harzburgites, an ophiolitic mélange zone, and the Moho consisting of dunites and harzburgite xenoliths (Xiong et al., 2016, 2022; Zhang et al., 2016a, 2020b, 2024b). The magmatic crust in these ophiolites has a variable thickness and commonly outcrops as blocks within the mélange (Zhang et al., 2016a, 2024b).

The formation time of the YTO has been extensively studied using high-precision zircon U-Pb dating (Table S1). According to these age data, the YTO along the >1000-km-long suture formed in a relatively short time span of ~130–120 Ma (Fig. 1c; Table S1). Notably, there is strong consistency in the age results obtained across different lithologies, from the upper crust on the top, through the lower crustal gabbros, to the mafic dikes that crosscut the mantle (Fig. 1c) (see Liu et al., 2016). The formation time of the ophiolite aligns closely, within error, with that of the underlying metamorphic sole (Duan et al., 2022; Guilmette et al., 2009, 2012; Zhang et al., 2016a, 2019, 2023b) and the overlying radiolarian cherts (Ziabrev et al., 2003) (Fig. 1c). The origin for such rapid accretion of the oceanic crust has been discussed previously. The proposed models are diverse, for example, subduction initiation and forearc spreading at the Asian continental margin (Dai et al., 2013, 2021), back-arc spreading in response to the intra-oceanic subduction of the Neo-Tethys (Guilmette et al., 2009, 2012; Hébert et al., 2012), detachment faulting at an ultraslow-spreading ridge (Liu et al., 2016), and southward trench migration and slab rollback (Xiong et al., 2016). Nonetheless, a consensus has emerged that the Early Cretaceous YTO represents only a snapshot of the vast Neo-Tethys Ocean, which has a spreading history of approximately 120 Myrs (Liu et al., 2023).

Geochemistry of the Yarlung-Tsangpo ophiolites

A series of studies have been undertaken to determine the geochemical characteristics of the YTO. Here we compiled available bulk-rock and mineral data for basalts and diabases ($n = 379$) and lower crustal gabbros ($n = 121$) from the well-studied Xigaze ophiolite in the central YTS, as well as for mantle peridotites (harzburgite and lherzolite, $n = 550$) along the entire suture. These data, along with Re-Os isotopes and highly siderophile elements (HSE) of the mantle peridotites, will be utilized to discuss the tectonic setting, crustal accretion, and mantle dynamics of the YTO.

Based on the dataset, we examined the fundamental characteristics of the crustal rocks and mantle peridotites of the YTO. Basalts and diabases typically exhibit rare earth element (REE) patterns characterized by a weak depletion in light REE (LREE) and relatively flat middle and heavy REE (MREE and HREE). These REE patterns are similar to that of normal mid-ocean ridge basalt (N-MORB); some diabases and minor basalts, however, display marked REE depletions relative to N-MORB (Fig. 2a). Compared to N-MORB, the basalts and diabases of the YTO exhibit noticeable depletions in Nb and Ta, along with a subtle depletion in Ti. Additionally, some of these rocks are enriched in large ion lithophile elements (LILE), such as Rb and Ba (Fig. 2a). In the Th/Yb vs. Nb/Yb diagram, the basalts and diabases have large variations (Fig. 2b), with some samples having elevated Th/Yb ratios that plot within the field of oceanic arcs.

Lithologically, the lower crustal cumulate rocks of the YTO comprise olivine-rich troctolite, troctolite, olivine gabbro, layered gabbro, isotropic gabbro, and minor gabbro-norite and oxide gabbro (Girardeau et al., 1985; Liu et al., 2018b, 2021a, b). These rocks overall display significant compositional variations, indicating variably evolved characteristics from relatively primitive olivine-bearing rocks to more evolved olivine-free gabbros (Liu et al., 2018b). In the bulk Al_2O_3 vs. MgO correlation, the YTO cumulate rocks follow a fractionation trend that suggests an earlier precipitation of plagioclase compared to clinopyroxene (Fig. 2c). This trend is further exemplified by a rapid decrease in bulk Ca# [$100 \times \text{Ca} / (\text{Ca} + \text{Na})$], followed by a decline in bulk Mg# [$100 \times \text{Mg} / (\text{Mg} + \text{Fe}^{2+})$] (Fig. 2d). This pattern is consistent with a significant removal of Ca during the fractionation of plagioclase. Overall, the composition of the YTO cumulate rocks is comparable to that of abyssal cumulates at modern mid-ocean ridges and can be distinguished from SSZ cumulates (Fig. 2c, d; compiled by Zhang et al., 2022).

Similar to the lower crustal cumulate rocks, the mantle peridotites along the entire suture exhibit significant variation in their geochemical compositions, ranging from refractory to relatively fertile characteristics (Fig. 2e-g). Most of these samples are compositionally comparable to abyssal peridotites (Fig. 2e-g; data from the PetDB database: <https://search.earthchem.org/>). A small subset, however, displays relatively lower bulk and pyroxene Al_2O_3 , along with higher olivine forsterite [Fo; $100 \times \text{Mg} / (\text{Mg} + \text{Fe}^{2+})$] and spinel Cr# [$\text{Cr} / (\text{Cr} + \text{Al})$], resembling refractory forearc peridotites from the West Pacific subduction systems (Fig. 2e-g; PetDB database). The extensive compositional variations observed in the mantle peridotites of the YTO have been attributed to complex processes involving variable extents of mantle melting, melt-rock interaction, and prior melt depletions.

The geochemical characteristics of the YTO have been widely used to discriminate its tectonic setting. The enrichment in LILE and the depletion in certain high field strength elements (such as

Nb, Ta, Ti) in the basalts and diabases are regarded as diagnostic features supporting a SSZ origin for the YTO, either a forearc (Fig. 2a, b) (Dai et al., 2013, 2021; Malpas et al., 2003; Xiong et al., 2016) or back-arcs (Bédard et al., 2009; Hébert et al., 2012). Similarly, the refractory mantle peridotites, characterized by high spinel Cr# and elevated whole-rock LREE, have been interpreted as forearc or back-arc peridotites that interacted with slab-derived melts (Fig. 2e-g) (Bezard et al., 2011; Dubois-Côté et al., 2005; Dupuis et al., 2005; Xia et al., 2003; Xiong et al., 2016, 2017a, b; Xu et al., 2020). The arc-like geochemical features, however, are not present within the lower crustal cumulates. The gabbros in the YTO are characterized by an earlier crystallization of plagioclase relative to clinopyroxene, as evidenced by their geochemical compositions (Fig. 2c, d) and petrographic observations (Liu et al., 2018b, 2021b). This is consistent with the low-pressure crystallization of dry basaltic magmas (Grove et al., 1992). The presence of troctolite within the lower crustal cumulates of the YTO further support an anhydrous line of descent (Liu et al., 2018b). Moreover, intra-oceanic arcs associated with the subduction of the Neo-Tethyan slab are also absent along the suture (Wu et al., 2014; Zhang et al., 2014). To resolve the ongoing debate regarding the tectonic setting of the YTO, a clear distinction must be made. In the following discussions, we will illustrate how the understanding of the origin and tectonic setting of the YTO has undergone a paradigm shift and how ophiolite conundrum can be reconciled through a comprehensive comparison with the structures and geochemical characteristics of oceanic lithospheres at modern ultraslow-spreading ridges.

DISCUSSION

Thin and highly variable oceanic crusts

One of the most conspicuous geological features of the YTO is its extremely thin oceanic crusts along the entire suture. The entire belt of the YTO extends roughly in an E–W direction, the thickness of its magmatic crusts thus was measured in the field perpendicularly to this direction (Girardeau et al., 1985b; Liu et al., 2021a, b; Nicolas et al., 1981). In localities where the magmatic crust is well developed, the thickness of the crust also refers to the orientation of crust–mantle boundary and foliation of gabbros and mantle peridotites (Girardeau et al., 1985b; Liu et al., 2021a, b; Nicolas et al., 1981).

In the western segment, the magmatic crust is nearly absent; in the eastern segment, it is only minor; and in the central segment, it does not exceed 2–3 km in thickness (Fig. 3). This is markedly distinct from the Penrose-type oceanic lithosphere, which typically has a crustal thickness of 6–7 km, such as the East Pacific Rise and the Oman and Troodos ophiolites (Fig. 3). A thin crust may

have resulted from extensive tectonic dismemberment during ophiolite obduction and the subsequent Indo–Asian collision. However, field observations from previous studies (Girardeau et al., 1985b; Nicolas et al., 1981; Wang et al., 1987) indicate that this tectonic dismemberment is negligible, a finding that has also been confirmed by our field work (Liu et al., 2021a, b). Notably, an intrusive contact has been identified between the lower crustal gabbros and underlying mantle peridotites (Fig. 4a; Liu et al., 2016). In this intrusive contact, the peridotites are moderately serpentinized, while the gabbros are leucogabbros with abundant plagioclases (Fig. 4a). These features indicate a cold lithospheric mantle intruded by relatively evolved gabbros at shallow horizons. Additionally, pillow lavas on top of the crust are directly overlain by radiolarian cherts (Fig. 4b; Wang et al., 2017). The preservation of these primary contacts indicates that the internal structures of the YTO are relatively intact, suggesting that the thin crust is an inherent feature rather than a product of significant tectonic dismemberment.

The magmatic crust of the YTO is not only thin but also highly variable in both thickness and structure. Notably, the lower crust, defined as seismic layer 3, is discontinuous. For example, lower crustal gabbros are developed only in three localities (i.e., Jiding, Baigang, and Dazhuqu) within the Xigaze ophiolite of the central YTO (Girardeau et al., 1985a, b; Liu et al., 2016; Nicolas et al., 1981; Wang et al., 1987; Wu et al., 2014). In Baigang, the thickness of these gabbros does not exceed ~800 m, while in Jiding, it is only about ~350 m (Liu et al., 2016). The lower crustal gabbros, along with the mantle peridotites, are ubiquitously intruded by a second generation of mafic rocks. These later intrusions crosscutting the mantle unit consist of diabase, gabbro, rodingite, and foliated amphibolite, while those intruding the lower crustal unit are predominantly diabases (Fig. 5). In both cases, the proportion of these isolated mafic rocks increase upwards, transferring gradually into a sheeted diabase unit composed entirely of diabases. Overall, the isolated mafic rocks throughout the ophiolite sequence have variable orientations (Liu et al., 2021a). They commonly strike sub-parallelly to the high-temperature ductile deformation of mantle peridotites and the layering of lower crustal gabbros, thus having been termed “diabase sills” (Fig. 5; Girardeau et al., 1985; Liu et al., 2021a, 2022b; Nicolas et al., 1981; Wu et al., 2014). Diabases that are oblique or sub-perpendicular to the mantle and gabbros are also present (Fig. 5d). The sheeted diabase unit is dominated by diabase sills and has been defined as “sheeted sill complex” (Fig. 5f; Girardeau et al., 1985; Liu et al., 2021a, 2022b; Nicolas et al., 1981; Wu et al., 2014).

The along-axis crustal thickness in a W–E direction has been measured for the well-studied Xigaze ophiolite, from Jiding in the west to Dazhuqu in the east (Fig. 6). The result underscores a highly variable magmatic crust within the Xigaze ophiolite (Fig. 6b). In the west (Jiding) and east (Baigang–Dazhuqu), the crustal unit is relatively thick, with lower crustal gabbros of 350–800 m

and an upper crust no more than 2 km. These areas correspond to volcanic segments ~25 km long in the west and ~38 km long in the east along the paleo-ridge. In contrast, the central part of the Xigaze ophiolite, spanning about ~53 km, exhibits an extremely thin magmatic crust, which is even absent in some areas, resembling an amagmatic zone. Notably, lower crustal gabbros are also lacking in this zone. This highly variable and unevenly distributed oceanic crust suggests strong magma focusing during the formation of the Xigaze ophiolite (Liu et al., 2021b). The observation of thin or missing crust in the central amagmatic zone imply extensive exposure of the mantle on the ancient seafloor, bordered by smaller-scale volcanic segments (Fig. 6b). These features mimic the crustal architecture of modern ultraslow-spreading ocean ridges, which are characterized by highly focused magmatic accretion and mantle exposures (Cannat et al., 2003; Dick, 1989; Dick et al., 2003; Li et al., 2015; Mendel et al., 1997; Rommevaux-Jestin et al., 1997; Whitehead et al., 1984; Zhang et al., 2024a). This crustal architecture is distinctly different from the uniform and continuous oceanic crust typical of fast-spreading ridges (Babcock et al., 1998; Sinton and Detrick, 1992) and can also be distinguished from the relatively weak magma focusing and more closely spaced smaller volcanic centers characteristic of slow-spreading ridges (Lin and Morgan, 1992; Lin et al., 1990; Mendel et al., 1997; Rommevaux-Jestin et al., 1997; Tolstoy et al., 1993).

Episodic lower crustal accretion

A ~350-m-thick lower crust has been measured in the Jiding section of the Xigaze ophiolite to examine its compositional variation (Fig. 6a, 7a; Liu et al., 2021a, b). The Jiding lower crust is composed of gabbroic rocks and diabase dikes/sills. The gabbroic rocks are predominately olivine-free, with only negligible occurrence of olivine gabbros and gabbronorite. They are mainly layered gabbros with evident magmatic foliations, which are defined by shape preferred orientations of clinopyroxene and plagioclase. The upper 20–30 m of the section is made up of oxide gabbros. It is evident that the Jiding lower crust is extremely thin with respect to the ~2.5-km-thick Atlantis Bank gabbro massif at the Southwest Indian Ridge (Dick et al., 2019; Zhang et al., 2020c) and the ~5-km-thick lower crust in the Oman ophiolite (Koepke et al., 2022; Müller et al., 2022) (Fig. 7b). It probably indicates a lower magma flux in Jiding with respect to Atlantis Bank and Oman (Liu et al., 2021b) or may be simply due to magmas for the Jiding lower crustal accretion were transported from elsewhere, as what happened at transform faults or mid-ocean ridges (Basch et al., 2024; Brunelli et al., 2020). It should be noted that primitive gabbroic rocks (troctolites and olivine gabbros) have been widely identified in other localities of the Xigaze ophiolite, i.e., Dazhuqu and Baigang (Liu et al., 2018b), which resemble the primitive gabbroic rocks of modern slow- and ultraslow-spreading ridges (Lissenberg and Dick, 2008; Sanfilippo et al., 2018; Zhang et al., 2021). It indicates that the

Jiding lower crust was more evolved than those in Dazhuqu and Baigang (Liu et al., 2018b). However, even considering the entire Xigaze ophiolite together, its lower crust is still thin and does not exceed 800 m in thickness (Liu et al., 2018b). Therefore, it is more reasonable that the Xigaze ophiolite has a lower magma flux than Atlantis Bank and Oman.

The cryptic chemical variation of the Jiding lower crust is also distinct from those observed at Atlantis Bank and Oman. Based on our previous geochemical data, the Jiding gabbros display episodic chemical variations in both bulk-rock and mineral compositions (Fig. 7c-f), which can be categorized into five units. Within individual units, clinopyroxene Mg#, Cr₂O₃, and bulk Mg# gradually decrease upwards, while clinopyroxene TiO₂ increases. There are clear breaks at unit boundaries. Such features were interpreted by that the lower oceanic crust is constructed by several magma intrusions (Dick et al., 2000, 2002). In each intrusion, interstitial evolved melts in the crystal mush can be transferred towards the top of the sequence by compaction, as evidenced by the pervasive occurrence of reactive porous flow throughout the lower crust (e.g., Boulanger et al., 2021; Lissenberg and MacLeod, 2016; Sanfilippo et al., 2020). Such episodic lower crustal accretion is not exclusive to the Jiding section, but has also been identified at Atlantic Bank, as evidenced by the downhole variations recorded in ODP/IODP Hole 735B and U1473A (Dick et al., 2000, 2002; Zhang et al., 2020c). A clear distinction, however, exist between Atlantis Bank and Jiding, particularly in the size of individual intrusive cycles. The intrusive cycles at Atlantis Bank are nearly an order of magnitude larger than those at Jiding (~500 m in Atlantis Bank compared to ~100 m in Jiding; Fig. 7b). This difference can be attributed to variations in their magma fluxes, with Atlantis Bank representing an anomalous, moderate magma budget compared to the rest of the ultraslow-spreading Southwest Indian Ridge (Dick et al., 2019; Zhou and Dick, 2013).

Episodic lower oceanic crustal accretion is not typical of the supposed oceanic crust of the Oman ophiolite (Coogan, 2014; Koepke et al., 2022; MacLeod and Yaouancq, 2000; Müller et al., 2022) nor of modern fast-spreading oceanic crust, such as those at the East Pacific Rise (Coogan, 2014; Lissenberg et al., 2013). The geochemical compositions of the Oman lower crust are relatively homogeneous throughout the majority of the section, with the exception of the uppermost horizon, which contains numerous varitextured gabbros representing the fossil axial melt lens (Fig. 7b; MacLeod and Yaouancq, 2000). Recent studies from the Oman drilling project have identified noticeable compositional variations between the upper and lower foliated gabbro units (see Fig. 7b), which were interpreted by hybrid accretion with the contributions from the axial melt lens on the top and the gabbro sills underneath (Koepke et al., 2022; Müller et al., 2022). Nonetheless, the Oman ophiolite is characterized by a much higher magma supply supporting ~5-km-thick lower

oceanic crust. The Oman ophiolite thus represents a higher class of magma budget compared to Jiding (low) and Atlantis Bank (moderate) (Fig. 7b; Liu et al., 2021b).

Recognition of detachment fault and its evolution

The thin and highly variable magmatic crust along the entire suture, coupled with episodic lower crustal accretion in the central segment, suggests that the YTO formed under a condition of low magma supply, likely at an ancient ultraslow-spreading ridge (Liu et al., 2021b). This argument is further substantiated by the occurrence of detachment faults and oceanic core complexes identified within the YTO. Such interpretations are supported by multidisciplinary evidence encompassing structural geology, petrology, geochronology, geochemistry, and paleomagnetism.

It has been suggested that the mantle unit of the Xigaze ophiolite has undergone large-scale internal rotations, revealed by paleomagnetic studies of crosscutting diabase dikes (Maffione et al., 2015). This behavior is analogous to the block rotations observed in active detachment faults of modern mid-ocean ridges (MacLeod et al., 2011; Morris et al., 2009). Structural studies have provided further evidence for the existence of detachment faults. (1) Retrograde serpentinite fault zones were identified in both the Xigaze ophiolite in the central segment (Li et al., 2016, 2021) and the Zedong ophiolite in the eastern segment (Li et al., 2025b), where talc-rich schists replaced the primary ductile deformation of mantle peridotites. (2) Tectono-sedimentary breccias are developed on the top of those serpentinite fault zones, consisting of serpentinite breccias in the sheared serpentinite matrix (Li et al., 2025b). These structural features resemble those within the interface of detachment faults at both modern slow- and ultraslow-spreading ridges (Escartín et al., 2003; Hansen et al., 2013; MacLeod et al., 2002) and the Apline-Apennine ophiolites (e.g., Manatschal et al., 2011).

Numerous mafic dikes/sills crosscut the serpentinitized mantle peridotites of the YTO. They are composed mainly of diabase and gabbro, some of which are subject to extensive hydrothermal alteration and ductile deformation, resulting in the formation of rodingite and foliated amphibolite, respectively (Fig. 5a-c; Li et al., 2016; Liu et al., 2021a; Zhang et al., 2016a, 2023c; Zhou et al., 2023). This is a notable feature, as diabases in oceanic lithospheres are typically generated at shallow depths and are not expected to exist in the mantle. The geological and structural characteristics of the YTO suggest that mantle exhumation may have occurred prior to the intrusion of fresh diabase dikes (Liu et al., 2016). Rodingites are typically regarded as alteration products of mafic dikes, formed through hydrothermal alteration by Si- and Ca-rich fluids during serpentinitization (Bach and Klein, 2009; Li et al., 2007). The widespread presence of rodingites

within the serpentinized mantle peridotites thus indicates that some amounts of magmas have intruded into the mantle before it was exhumed and serpentinized (Liu et al., 2016, 2022b; Wu et al., 2014). Additionally, petrological studies reveal that the foliated amphibolites within the YTO mantle peridotites underwent relatively high-temperature metamorphism under the amphibolite-facies condition (600–900 °C, 3kbar), suggesting ductile deformation of their protoliths (Liu et al., 2021a; Zhang et al., 2023c; Zhou et al., 2023). As noticed above, all the mafic rocks within the YTO share consistent formation ages of 130–120 Ma. In individual ophiolite sections, the temporal duration for the generation of the mafic rocks therein is even shorter (i.e., < 5 Myr; Liu et al., 2016). Notably, the diabase and gabbro dikes in the mantle, along with the rodingite and foliated amphibolite, are geochronologically indistinguishable from the oceanic crust (Fig. 1c). This indicates a dynamic linkage between melt crystallization and mantle exhumation, accompanied by hydrothermal alteration (serpentinization and rodingitization), ductile deformation, and high-temperature metamorphism (foliated amphibolites). These findings align well with previous paleomagnetic and structural studies that support the occurrence of mantle exhumation, rotation, and serpentinite ductile shear zones in the YTO (Li et al., 2016, 2025b; Maffione et al., 2015).

All lines of evidence from petrology, geochronology, geochemistry, paleomagnetism, and structural geology confirm the occurrence of detachment faults in the YTO (Li et al., 2016, 2025b; Liu et al., 2014, 2016, 2021a, 2021b; Maffione et al., 2015). This conclusion is further supported by a recent microstructural study revealing a multi-stage deformation history for the YTO mantle peridotites, which encompasses high-temperature ductile shearing followed by the development of detachment-fault-hosted mylonitization (Zhang et al., 2025). The formation and evolution of these detachment faults, in particular the associated footwall rotations, are the primary mechanisms to account for the pervasive occurrence of variably oriented mafic dikes in both the mantle and lower crust (Liu et al., 2021a). Similar mafic dikes are also not uncommon in other slow-spreading-type ophiolites and in boreholes from ocean drillings at modern slow- and ultraslow-spreading ridges (Liu et al., 2024 and references therein). The geometric structures of these mafic dikes, along with their relationships with the host mantle and lower crust, exhibit significant diversity (e.g., Blackman et al., 2006; Godard et al., 2009). It can be explained by multi-stage dike intrusions into the variably exhumed and rotated mantle peridotites and gabbros at the detachment footwall (Liu et al., 2021a). This complexity contributes to an intricate interplay between magma injection, mantle exhumation, and rotation.

In summary, the thin and highly variable crust, episodic lower crustal accretion, and occurrence of detachment faults combined lead us to conclude that the YTO formed under a low magma budget at an ancient ultraslow-spreading ridge. This ridge exhibited along-axis magma

focusing (Fig. 8a) and cross-axis detachment faulting (Fig. 8b). The YTO is clearly distinguishable from the Penrose model, which describes a layered oceanic crust of ~6–7 km thick. Instead, it more resembles the Chapman-type oceanic lithosphere characterized by thin or absent oceanic crusts, along with the development of detachment faults and oceanic core complexes (Escartín and Canales, 2011; Maffione et al., 2013). In fact, Chapman-type ophiolites that share similar crustal architectures with the YTO are significantly more prevalent than Penrose-type ophiolites, with an estimated ratio of 90% vs. 10%, particularly along the extensive Neo-Tethyan ophiolite belt stretching from the Alps to Southeast Asia (Fig. 3; Liu et al., 2024). This suggests that either Chapman-type oceanic lithospheres have a greater likelihood of preservation compared to Penrose-type (e.g., Ganade et al., 2024) or that slow- and ultraslow-spreading ridges were more abundant than fast-spreading ridges throughout the evolutionary history of the Neo-Tethys Ocean.

The highly heterogeneous mantle underneath

The thin crust of the YTO demonstrates an overall low magma supply along the paleo-ridge. It implies a relatively fertile residual mantle along the entire suture, which however is not the case for the mantle peridotites underneath. In Fig. 9, we reconstructed the compositional variations of mantle peridotites along the >1300-km-long Neo-Tethyan ocean ridge represented by the YTO. The results indicate a significant variation in mantle composition, such as bulk Al₂O₃, spinel Cr#, and clinopyroxene Yb (Fig. 9b-d). This is further exemplified by considerable variations in olivine Fo and pyroxene Al₂O₃ (not shown here). Such large variations are evident not only along the entire suture, but also within individual segments (i.e., western, central, and eastern) and even in single ophiolite sections. Based on spinel Cr#, we calculated the along-axis variation in the extent of mantle melting using the empirical formula: $F = 9 \times \ln(\text{Cr\#}) + 23$ (F means melting extent; Warren, 2016), which also demonstrates dramatic changes (Fig. 9e). Assuming that the highest values of F (17.8%–20.1%) for the mantle peridotites in each ophiolite section represents the maximum extent of melting in the 60-km deep melting regime, a simple mass balance calculation yields an estimated oceanic crustal thickness that is as high as 6.0–6.8 km. This estimation, however, does not align with the observed crustal thickness for the YTO, where the magmatic crust is absent in its western and eastern segments and does not exceed 2–3 km in its central segment. Considering that the influence of tectonic dismemberment for the crustal thickness of the YTO is insignificant, mantle melting beneath the ancient Neo-Tethyan mid-ocean ridge should be limited.

An explanation for the discrepancy between a refractory mantle and a thin crust is that the mantle has undergone a complex melting history rather than a single-stage sub-ridge melting (Liu et al., 2019; Xu et al., 2020; Zhang et al., 2017, 2023a; Zhu et al., 2025), a scenario that is consistent

with the ubiquitous ultra-depleted mantle domains beneath the Mid-Atlantic and Gakkell Ridges revealed by Hf–Nd isotopes (e.g., Sanfilippo et al., 2024; Sani et al., 2023; Stracke et al., 2011, 2019). This argument is supported by the extremely depleted compositions in Os isotopes along the YTO mantle (Fig. 9f), which confirm that these mantle peridotites have experienced ancient melting (up to ~2.2 Ga) prior to their entry into the sub-ridge melting regime (Gong et al., 2016; Lian et al., 2024; Liu et al., 2012, 2019; Niu et al., 2015; Xu et al., 2020, 2025; Zhang et al., 2020a, b, 2023a; Zhu et al., 2025). The evidence for multi-stage mantle melting is further illustrated by the correlations between the $^{187}\text{Os}/^{188}\text{Os}$ ratios vs. the $^{187}\text{Re}/^{188}\text{Os}$ ratios and bulk Al_2O_3 contents (Fig. 10a, b). It is on one hand because Re–Os isotope system is robust against post-melting metasomatism, thus could retain its long-term depletions (Liu et al., 2008; Rudnick and Walker, 2009). On the other hand, as a long-lived isotope system, osmium isotopic ratios remain nearly constant during recent melting events, while lithophile fertility indexes can change dramatically (Lassiter et al., 2014). In the correlation between $^{187}\text{Os}/^{188}\text{Os}$ and bulk Al_2O_3 , the mantle peridotites in the YTO exhibit roughly three linear trends, with slopes becoming steeper from high to low Al_2O_3 contents (Fig. 10b). This feature indicates a shift in the Al–Os isochron, which is not uncommon for abyssal peridotites (Lassiter et al., 2014) and other ophiolitic mantle peridotites (Li et al., 2025a; Wang et al., 2023). The Al–Os isochron shift can be attributed to the multi-stage melting experienced by mantle peridotites, significantly affecting Al without similarly impacting Os. For abyssal peridotites, this shift is thought to be caused by recent melting beneath the mid-ocean ridge that overprints prior melt depletion events (Lassiter et al., 2014). A similar conclusion of multi-stage mantle melting has also been achieved by studies on Hf–Nd isotopes of MORB and abyssal peridotites (e.g., Sanfilippo et al., 2024; Sani et al., 2023; Stracke et al., 2011, 2019). In the case of ophiolitic peridotites, the melting history may involve more complex processes, including ancient melting, sub-ridge melting, and hydrous melting in a SSZ setting (Li et al., 2025a; Wang et al., 2023). In the context of tectonic evolution of the Neo-Tethys Ocean, we tentatively propose three stages of mantle melting events (Fig. 10b), that is, ancient melting dating back to ~2.2 Ga (the oldest T_{RD} age for the YTO peridotites; Xu et al., 2020), sub-ridge melting at the Neo-Tethyan mid-ocean ridge at ~130 Ma (the formation age of the YTO), and a rapid overprinting by hydrous melting in a forearc setting at ~125 Ma. This multi-stage mantle melting model can effectively explain the large compositional variations in the YTO mantle peridotites, which overlap those of abyssal and forearc peridotites (Fig. 2e-g; Fig. 10a-b).

HSE compositions provide additional insights into the melting history of mantle peridotites in the YTO. Due to their extremely high metal/silicate partition coefficients, HSE in the mantle, at least after the so-called “late-veener hypothesis”, are preferentially preserved in sulfides. In a

simplified scenario, HSE in mantle peridotites might have significant fractionations via high extents (>15%–20%) of mantle melting or intense sulfide-rich melt percolation, which can change in the behavior of sulfides under these conditions (Büchl et al., 2002; Liu et al., 2009; Lorand et al., 1999; Luguet and Reisberg, 2016; Xu and Liu, 2019). The HSE compositions of most YTO mantle peridotites are chondritic and comparable to that of the primitive upper mantle (PUM; Becker et al., 2006) (Fig. 10c, d). In contrast, some samples exhibit fractionated HSE compositions, displaying variable Re and depleted Pt and Pd with respect to the PUM (Fig. 10c). Meanwhile, the YTO mantle peridotites generally have PUM-like $(\text{Pt}/\text{Pd})_N$ ratios, while the most refractory peridotites have elevated $(\text{Pt}/\text{Pd})_N$ ratio. Such elevated $(\text{Pt}/\text{Pd})_N$ ratios, commonly documented in the sub-arc mantle (including forearc peridotites and mantle wedge xenoliths), are attributed to the formation of Pt-Fe alloys during multi-stage mantle melting (Kepezhinskas and Defant, 2001; Xu and Liu, 2019). Although the $(\text{Pt}/\text{Pd})_N$ ratios in the refractory YTO mantle peridotites is not as high as those in the sub-arc mantle, it implies that these samples must have experienced higher extents of mantle melting compared to other samples. Therefore, the HSE compositions bolster our argument based on Os isotopes, indicating that the YTO mantle peridotites are products of multi-stage mantle melting and reflecting a highly heterogeneous mantle beneath the Neo-Tethyan mid-ocean ridge.

Ancient mantle domains responsible for the heterogeneity of Neo-Tethyan asthenosphere have been thoroughly explained as resulting from the recycling of continental (Gong et al., 2016; Griffin et al., 2016), sub-oceanic (Liu et al., 2019; Zhang et al., 2020a, b; Zhang et al., 2023a), or subduction-modified peridotites (Liu et al., 2012). Similar hypotheses have also been proposed for the mantle peridotites along the entire Neo-Tethyan ophiolite belt (Liu et al., 2021c and references therein). The occurrence of portion of recycled mantle in the asthenosphere is a common feature at modern slow- and ultraslow-spreading ridges, where abyssal peridotites have been dated to >2.5 Ga (Harvey et al., 2006; Lin et al., 2024; Liu et al., 2008, 2022a; Sanfilippo et al., 2024; Sani et al., 2023; Stracke et al., 2011, 2019). This would result in a common phenomenon that thin oceanic crust is supported by a highly heterogeneous mantle (D'Errico et al., 2016; Zhou and Dick, 2013), highlighting the need for a careful examination of the mantle's nature to better understand the crust-mantle relations.

Thin crust supported by a highly heterogeneous mantle is also a typical feature in modern and fossil magma-poor OCT settings, such as the Iberia-Newfoundland in the North Atlantic Ocean and the Alpine Tethys (e.g., Manatschal et al., 2011; Müntener and Manatschal, 2006; Rampone and Hofmann, 2012; Rampone and Sanfilippo, 2021). Notably, the Alpine-Apennine ophiolites are suggested to contain abundant SCLM that were stagnated within the asthenosphere during continental rifting (McCarthy and Müntener, 2015; Rampone and Hofmann, 2012; Rampone and

Sanfilippo, 2021). These features are quite similar to those of the YTO. However, there are fundamental distinctions between the Yarlung-Tsangpo Neo-Tethys Ocean and the Alpine Tethys Ocean (i.e., the Piemonte-Ligurian Ocean). The Yarlung-Tsangpo Neo-Tethys was a mature open ocean that are thousands-of-kilometers wide at the formation time of the YTO, that is, ~130–120 Ma, as evidenced by paleomagnetic data for the Indian and Asian continents (see Hu et al., 2016; Huang et al., 2024). The Yarlung-Tsangpo Neo-Tethys thus is different from the embryonic Piemonte-Ligurian Ocean but more resembles modern ultraslow-spreading open oceans, such as the Southwest Indian Ridge. Accordingly, the YTO should have formed in an ultraslow-spreading ridge, rather than an OCT setting like the Alpine-Apennine ophiolites.

Paradigm shift in understanding the YTO: new models for ophiolite conundrum

The origin of ultraslow-spreading ridges for the YTO and the identification of ancient mantle domains therein provide new insights into the tectonic setting for its formation. In the context of ultraslow seafloor spreading, the observed arc-like features can alternatively be attributed to two new models, rather than the subduction initiation rule typically applied to global SSZ-type ophiolites (Stern, 2004; Whattam and Stern, 2011).

One explanation for the arc-like geochemical signatures in the YTO involves the melting of ancient mantle domains within a heterogeneous asthenosphere under the ultraslow-spreading Neo-Tethyan mid-ocean ridge (Fig. 11a). These ancient peridotites are likely recycled sub-continental lithospheric mantle (SCLM), arc mantle, or ancient sub-oceanic mantle (e.g., Liu et al., 2021c). This model is supported by the occurrence of anomalous compositions documented in MORBs at modern slow- and ultraslow-spreading ridges. For instance, a refertilized SCLM is thought to have the potential to generate the Dupal isotope anomaly in MORBs observed in both the Southern Hemisphere (Kempton et al., 2002; Mahoney et al., 1992) and the Gakkel Ridge, Arctic Ocean (Goldstein et al., 2008). Moreover, the SCLM has long been recognized as a candidate source for generating enriched MORBs distant from mantle plumes (McKenzie and O'Nions, 1983; Yang et al., 2020), as well as the alkaline, K-rich basalts at the Southwest Indian Ridge and Mid-Atlantic Ridge (Nauret et al., 2011; Wang et al., 2020). Recycled arc mantle, on the other hand, is believed to be prevalent by Urann et al. (2020), based on a handful of peridotites in a Mid-Atlantic Ridge section. The widespread contamination of the asthenosphere by these arc mantle can result in a hemisphere-scale upper mantle heterogeneity, generating BABB (back-arc basin basalt)-like MORBs frequently encountered at global ocean ridges barring the Pacific Ocean as inferred for instance by Yang et al. (2021). It should be noted that these findings from modern ocean ridges do not imply that the pervasive occurrence of arc-like lavas in global ophiolites can be well accounted

for by the melting of recycled mantle in the asthenosphere. Nonetheless, it provides an additional perspective for studying certain ophiolites, suggesting that caution should be exercised when using basalts to discriminate the tectonic setting of ophiolites (Liu et al., 2021c).

The ophiolite conundrum of the YTO can also be elucidated by a subduction jump and re-initiation model proposed by recent studies on the Yarlung-Tsangpo metamorphic soles (Zhang et al., 2019). Metamorphic soles associated with ophiolites are widely regarded as geological indicators for subduction initiation (e.g., Wakabayashi and Dilek, 2003). It is shown that metamorphic soles along the YTS are compositionally similar to the magmatic crust of the YTO (e.g., Zhang et al., 2019). Their protolith and metamorphic ages are also coeval with the formation ages of ophiolites (Fig. 1c; Duan et al., 2022; Guilmette et al., 2009, 2012; Zhang et al., 2019, 2023b). These features thus suggest a rapid tectonic transition from an ultraslow-spreading ridge to a subduction zone (Duan et al., 2022; Liu et al., 2022b; Zhang et al., 2019, 2023b). A notable feature of slow- and ultraslow-spreading ridges is the highly heterogeneous crustal structures on the seafloor. It has been established that detachment and transform faults both can lead to significant exhumation of the mantle, forming topographic highs with respect to the ambient oceanic crust (Baines et al., 2003; Ligi et al., 2022), such as the Atlantis Bank, which was uplifted >3 km above adjacent seafloor (Baines et al., 2003). The subduction of such ocean ridges would result in a ridge-trench collision and the subsequent juxtaposition of oceanic core complexes at the continental margin (Fig. 11b; Liu et al., 2022b; Zhang et al., 2019). In such a scenario, a new subduction zone may re-initiate on the opposite side due to jump of the subduction front (Fig. 11b). Dehydration of the slab from this new subduction zone, along with the subsequent flux melting of the overlying mantle wedge (the fossil oceanic core complex), can generate arc-like magmatism superimposed on the pre-existing MORB-type oceanic crust. Such a model can also explain the rapid transition from mid-ocean ridge to subduction zone, during which the metamorphic soles were generated and had similar compositions and ages with the magmatic crusts of the YTO.

The aforementioned models provide a new perspective for resolving the long-standing debate regarding the tectonic setting of the YTO. Many geological and geochemical features of the YTO can be more effectively explained in the context of ultraslow seafloor spreading and its subsequent evolution, rather than simply relying on the previously accepted model of subduction initiation. This insight has the potential to lead to a paradigm shift in our understanding of this renowned, extensively studied ophiolite belt. Moreover, the YTO is not an isolated case, because slow- and ultraslow-spreading oceanic lithospheres, as described in the Chapman model, are likely to be prevalent within global ophiolites, particularly those associated with the Neo-Tethys extending from the Alps to Southeast Asia (Liu et al., 2024). Therefore, comparative studies with modern slow- and

ultraslow-spreading ridges should emerge as a new area of research, paving the way for a transformative shift in our knowledge of global ophiolites from the Penrose model to Chapman model.

CONCLUDING REMARKS

The understanding of ophiolites has been significantly reshaped since the pioneering work that innovatively linked ophiolites to island arcs (Miyashiro, 1973). This marked a paradigm shift, positioning ophiolites as geological records for subduction initiation, rather than products of seafloor spreading at mid-ocean ridges. If this hypothesis is valid, it means that the vast oceans that existed in Earth's geological history have completely subducted back into the deep mantle. However, the notion of a SSZ origin for ophiolites has been called into question, as it is suggested that MORB can display arc-like geochemical signatures. The key to resolving this issue lies in a comprehensive understanding of the underlying mantle. The extent and heterogeneity of mantle depletion serve as critical factors that modulate the thickness, composition, and characteristics of oceanic crusts. This knowledge, obtained from studies of modern slow- and ultraslow-spreading ridges, should be applied to ophiolite research, to identify those that may have formed under similar conditions. Ophiolites, in turn, are valuable supplements for the study of oceanic lithospheres at modern ocean ridges, as they typically provide continuous observations of lithology, composition, and structures of oceanic lithospheres.

ACKNOWLEDGEMENTS

We would like to express our gratitude to Riccardo Tribuzio for his invitation to contribute to this special issue celebrating the journal's 50th anniversary. We greatly acknowledge the editorial work by Editors Daniele Brunelli and Alessandra Montanini and the constructive comments from reviewers Gianreto Manatschal and Alessio Sanfilippo, which greatly improved the quality of the manuscript. This paper reviews the major findings of our studies on the Yarlung-Tsangpo ophiolites over the last decade. We acknowledge the contributions of Riccardo Tribuzio and Henry J. B. Dick to our research, as well as the assistances of Liang-Liang Zhang, Yin-Zheng Lin, Xiao-Ni Li, Zi-Hang Huang, Jia-Ning Zhu, Hong-Yu Li, De-Xiang Hong, Yong-Qiang Hu, and Liu-Yang Yang during the field work. Yang Chu is thanked for his help in the figure designation. Tong Liu acknowledges the support by the CAS Scholarship during his visit to Universität Münster, Germany, where this paper was completed. This study was financially supported by the National Science Fund for Distinguished Young Scholars (42025201), the Strategic Priority Program (Category B) of Chinese Academy of Sciences (CAS) (XDB0710000), and the Key Research Program of the Institute of Geology and Geophysics, CAS (IGGCAS-202202).

REFERENCES

- Anonymous, 1972. Penrose field conference on ophiolites. *Geotimes*, 17: 22-24.
- Ao S.J., Xiao W.J., Jafari M.K., Talebian M., Chen L., Wan B., Ji W.Q. and Zhang Z.Y., 2016. U–Pb zircon ages, field geology and geochemistry of the Kermanshah ophiolite (Iran): From continental rifting at 79 Ma to oceanic core complex at ca. 36 Ma in the southern Neo-Tethys. *Gondwana Research*, 31: 305-318. <https://doi.org/10.1016/j.gr.2015.01.014>.
- Babcock J.M., Harding A.J., Kent G.M. and Orcutt J.A., 1998. An examination of along-axis variation of magma chamber width and crustal structure on the East Pacific Rise between 13°30'N and 12°20'N. *Journal of Geophysical Research: Solid Earth*, 103: 30451-30467. <https://doi.org/10.1029/98JB01979>.
- Bach W. and Klein F., 2009. The petrology of seafloor rodingites: Insights from geochemical reaction path modeling. *Lithos*, 112: 103-117. <https://doi.org/10.1016/j.lithos.2008.10.022>.
- Baines A.G., Cheadle M.J., Dick H.J.B., Scheirer A.H., John B.E., Kuszniir N.J. and Matsumoto T., 2003. Mechanism for generating the anomalous uplift of oceanic core complexes: Atlantis Bank, southwest Indian Ridge. *Geology*, 31: 1105-1108. <https://doi.org/10.1130/G19829.1>.
- Basch V., Sanfilippo A., Snow J.E., Loocke M. and Zanetti A., 2024. Accretion of the Lower Oceanic Crust at Fast-Spreading Ridges: Insights from Hess Deep (East Pacific Rise, IODP Expedition 345). *Journal of Petrology*, 65: egae048. <https://doi.org/10.1093/petrology/egae048>.
- Becker H., Horan M.F., Walker R.J., Gao S., Lorand J.P. and Rudnick R.L., 2006. Highly siderophile element composition of the Earth's primitive upper mantle: Constraints from new data on peridotite massifs and xenoliths. *Geochimica et Cosmochimica Acta*, 70: 4528-4550. <https://doi.org/10.1016/j.gca.2006.06.004>.
- Bédard É., Hébert R., Guilmette C., Lesage G., Wang C.S. and Dostal J., 2009. Petrology and geochemistry of the Saga and Sangsang ophiolitic massifs, Yarlung Zangbo Suture Zone, Southern Tibet: Evidence for an arc–back-arc origin. *Lithos*, 113: 48-67. <https://doi.org/10.1016/j.lithos.2009.01.011>.
- Bezard R., Hébert R., Wang C., Dostal J., Dai J. and Zhong H., 2011. Petrology and geochemistry of the Xiugugabu ophiolitic massif, western Yarlung Zangbo suture zone, Tibet. *Lithos*, 125: 347-367. <https://doi.org/10.1016/j.lithos.2011.02.019>.
- Blackman D.K., Ildefonse B., John B.E., Ohara Y., Miller D.J. and MacLeod C.J., 2006. Proc. IODP, 304/305. College Station TX. <https://doi.org/10.2204/iodp.proc.304305.2006>. (Integrated Ocean Drilling Program Management International, Inc.).
- Brunelli D., Sanfilippo A., Bonatti E., Skolotnev S., Escartín J., Ligi M., Ballabio G. and Cipriani A., 2020. Origin of oceanic ferrodiorites by injection of nelsonitic melts in gabbros at the Vema Lithospheric Section, Mid Atlantic Ridge. *Lithos*, 368-369: 105589. <https://doi.org/10.1016/j.lithos.2020.105589>.
- Bortolotti V., Chiari M., Marcucci M., Marroni M., Pandolfi L., Principi G. and Saccani E., 2004. Comparison among the Albanian and Greek ophiolites: In search of constraints for the evolution of the Mesozoic Tethys Ocean. *Ophioliti*, 29: 19-35.
- Boulanger M., France L., Ferrando C., Ildefonse B., Ghosh B., Sanfilippo A., Liu C.Z., Morishita T., Koepke J. and Bruguier O., 2021. Magma-Mush Interactions in the Lower Oceanic Crust: Insights From Atlantis Bank Layered Series (Southwest Indian Ridge). *Journal of Geophysical Research Solid Earth*, 126: e2021JB022331.

<https://doi.org/10.1029/2021JB022331>.

- Büchl A., Brüggmann G., Batanova V.G., Münker C. and Hofmann A.W., 2002. Melt percolation monitored by Os isotopes and HSE abundances: a case study from the mantle section of the Troodos Ophiolite. *Earth and Planetary Science Letters*, 204: 385-402. [https://doi.org/10.1016/S0012-821X\(02\)00977-9](https://doi.org/10.1016/S0012-821X(02)00977-9).
- Cannat M., Rommevaux-Jestin C. and Fujimoto H., 2003. Melt supply variations to a magma-poor ultra-slow spreading ridge (Southwest Indian Ridge 61° to 69°E). *Geochemistry, Geophysics, Geosystems*, 4(8), 9104, doi:10.1029/2002GC000480. <https://doi.org/10.1029/2002GC000480>.
- Chang C.F. and Zheng X.L., 1973. Tectonic features of the Mount Jolmo Lungma region in southern Tibet, China. *Chinese Journal of Geology*, 8: 1-12 (in Chinese).
- Coogan L.A., 2014. 4.14 - The Lower Oceanic Crust. In: H.D. Holland and K.K. Turekian (Eds.). *Treatise on Geochemistry (Second Edition)*, Oxford, Elsevier: 497-541. <http://dx.doi.org/10.1016/B978-0-08-095975-7.00316-8>.
- D'Errico M.E., Warren J.M. and Godard M., 2016. Evidence for chemically heterogeneous Arctic mantle beneath the Gakkel Ridge. *Geochimica et Cosmochimica Acta*, 174: 291-312. <https://doi.org/10.1016/j.gca.2015.11.017>.
- Dai J.G., Wang C.S., Hébert R., Santosh M., Li Y.L. and Xu J.Y., 2011. Petrology and geochemistry of peridotites in the Zhongba ophiolite, Yarlung Zangbo Suture Zone: Implications for the Early Cretaceous intra-oceanic subduction zone within the Neo-Tethys. *Chemical Geology*, 288: 133-148. <https://doi.org/10.1016/j.chemgeo.2011.07.011>.
- Dai J.G., Wang C.S., Polat A., Santosh M., Li Y.L. and Ge Y., 2013. Rapid forearc spreading between 130 and 120Ma: Evidence from geochronology and geochemistry of the Xigaze ophiolite, southern Tibet. *Lithos*, 172-173: 1-16. <https://doi.org/10.1016/j.lithos.2013.03.011>.
- Dai J.G., Wang C.S., Stern R.J., Yang K. and Shen J., 2021. Forearc magmatic evolution during subduction initiation: Insights from an Early Cretaceous Tibetan ophiolite and comparison with the Izu-Bonin-Mariana forearc. *GSA Bulletin*, 133: 753-776. <https://doi.org/10.1130/B35644.1>.
- Dick H.J.B., 1989. Abyssal peridotites, very slow spreading ridges and ocean ridge magmatism. *Geological Society, London, Special Publications*, 42: 71-105. <https://doi.org/10.1144/GSL.SP.1989.042.01.06>.
- Dick H.J.B., Kvassnes A.J.S., Robinson P.T., MacLeod C.J. and Kinoshita H., 2019. The Atlantis Bank Gabbro Massif, Southwest Indian Ridge. *Progress in Earth and Planetary Science*, 6: 64. <https://doi.org/10.1186/s40645-019-0307-9>.
- Dick H.J.B., Lin J. and Schouten H., 2003. An ultraslow-spreading class of ocean ridge. *Nature*, 426: 405-412. <https://doi.org/10.1038/nature02128>.
- Dick H.J.B., Natland J.H., Alt J.C., Bach W., Bideau D., Gee J.S., Haggas S., Hertogen J.G.H., Hirth G., Holm P.M., Ildefonse B., Iturrino G.J., John B.E., Kelley D.S., Kikawa E., Kingdon A., LeRoux P.J., Maeda J., Meyer P.S., Miller D.J., Naslund H.R., Niu Y.L., Robinson P.T., Snow J., Stephen R.A., Trimby P.W., Worm H.U. and Yoshinobu A., 2000. A long in situ section of the lower ocean crust: results of ODP Leg 176 drilling at the Southwest Indian Ridge. *Earth and Planetary Science Letters*, 179: 31-51. [https://doi.org/10.1016/S0012-821X\(00\)00102-3](https://doi.org/10.1016/S0012-821X(00)00102-3).
- Dick H.J.B., Ozawa K., Meyer P.S., Niu Y., Robinson P.T., Constantin M., Hebert R., Maeda J., Natland J.H. and Hirth G., 2002. 10. Primary silicate mineral chemistry of a 1.5-km section of very slow spreading lower ocean crust: ODP hole 735B, Southwest Indian ridge. *Proceedings of Ocean Drilling Program, Scientific Results*, 176: 1-60.
- Dilek Y. and Furnes H., 2009. Structure and geochemistry of Tethyan ophiolites and their petrogenesis in subduction rollback systems. *Lithos*, 113: 1-20. <https://doi.org/10.1016/j.lithos.2009.04.022>.
- Dilek Y. and Furnes H., 2011. Ophiolite genesis and global tectonics: Geochemical and tectonic fingerprinting of ancient

- oceanic lithosphere. *Geological Society of America Bulletin*, 123, 387-411. <https://doi.org/10.1130/B30446.1>.
- Dilek Y. and Thy P., 2009. Island arc tholeiite to boninitic melt evolution of the Cretaceous Kizildag (Turkey) ophiolite: Model for multi-stage early arc-forearc magmatism in Tethyan subduction factories. *Lithos*, 113, 68-87. <https://doi.org/10.1016/j.lithos.2009.05.044>.
- Duan W.Y., Li X.P., Schertl H.P., Willner A.P. and Sun G.M., 2022. “Hot” subduction initiation and the origin of the Yarlung-Tsangbo ophiolites, southern Tibet: New insights from ultrahigh temperature metamorphic soles. *Earth and Planetary Science Letters*, 591: 117610. <https://doi.org/10.1016/j.epsl.2022.117610>.
- Dubois-Côté V., Hébert R., Dupuis C., Wang C.S., Li Y.L. and Dostal J., 2005. Petrological and geochemical evidence for the origin of the Yarlung Zangbo ophiolites, southern Tibet. *Chemical Geology*, 214: 265-286. <https://doi.org/10.1016/j.chemgeo.2004.10.004>.
- Dupuis C., Hébert R., Dubois-Côté V., Guilmette C., Wang C.S., Li Y.L. and Li Z.J., 2005. The Yarlung Zangbo Suture Zone ophiolitic mélange (southern Tibet): new insights from geochemistry of ultramafic rocks. *Journal of Asian Earth Sciences*, 25: 937-960. <https://doi.org/10.1016/j.jseaes.2004.09.004>.
- Escartín J. and Canales J.P., 2011. Detachments in Oceanic Lithosphere: Deformation, Magmatism, Fluid Flow, and Ecosystems. *Eos*, 92: 31. <https://doi.org/10.1029/2011EO040003>.
- Escartín J., Mével C., MacLeod C.J. and McCaig A.M., 2003. Constraints on deformation conditions and the origin of oceanic detachments: The Mid-Atlantic Ridge core complex at 15°45'N. *Geochemistry, Geophysics, Geosystems*, 4(8): 1067. <https://doi.org/10.1029/2002GC000472>.
- Fischer-Gödde M., Becker H. and Wombacher F., 2010. Rhodium, gold and other highly siderophile element abundances in chondritic meteorites. *Geochimica et Cosmochimica Acta*, 74: 356-379. <https://doi.org/10.1016/j.gca.2009.09.024>.
- Ganade C.E., Riel N., Manatschal G., Tesser L.R., Hermann J., Rubatto D., Weinberg R.F., Lanari P. and Kaus B.J.P., 2024. Exhumation of ultra-high pressure (UHP) rocks modulated by rifted margin-subduction feedback: Implications for their preservation in old collisional orogens. *Earth and Planetary Science Letters*, 643: 118893. <https://doi.org/10.1016/j.epsl.2024.118893>.
- Gansser A., 1964. *The Geology of the Himalayas*. Wiley-Interscience, New York, 1-289 pp.
- Girardeau J., Mercier J.C.C. and Wang X.B., 1985a. Petrology of the mafic rocks of the Xigaze ophiolite, Tibet. *Contributions to Mineralogy and Petrology*, 90: 309-321. <https://doi.org/10.1007/BF00384710>.
- Girardeau J., Mercier J.C.C. and Zao, Y.G., 1985b. Structure of the Xigaze Ophiolite, Yarlung Zangbo Suture Zone, southern Tibet, China: Genetic implications. *Tectonics*, 4: 267-288. <https://doi.org/10.1029/TC004i003p00267>.
- Godard M., Awaji S., Hansen H., Hellebrand E., Brunelli D., Johnson K., Yamasaki T., Maeda J., Abratis M., Christie D., Kato Y., Mariet C. and Rosner M., 2009. Geochemistry of a long in-situ section of intrusive slow-spread oceanic lithosphere: Results from IODP Site U1309 (Atlantis Massif, 30°N Mid-Atlantic-Ridge). *Earth and Planetary Science Letters*, 279: 110-122. <https://doi.org/10.1016/j.epsl.2008.12.034>.
- Goldstein S.L., Soffer G., Langmuir C.H., Lehnert K.A., Graham D.W. and Michael P.J., 2008. Origin of a ‘Southern Hemisphere’ geochemical signature in the Arctic upper mantle. *Nature*, 453: 89-93. <https://doi.org/10.1038/nature06919>.
- Gong X.H., Shi R.D., Griffin W., Huang Q.S., Xiong Q., Chen S.S., Zhang M. and O'Reilly S.Y., 2016. Recycling of ancient subduction-modified mantle domains in the Purang ophiolite (southwestern Tibet). *Lithos*, 262: 11-26. <https://doi.org/10.1016/j.lithos.2016.06.025>.
- Griffin W.L., Afonso J.C., Belousova E.A., Gain S.E., Gong X.H., González-Jiménez J.M., Howell D., Huang J.X., McGowan N., Pearson N.J., Satsukawa T., Shi R., Williams P., Xiong Q., Yang J.S., Zhang M. and O'Reilly S.Y.,

2016. Mantle Recycling: Transition Zone Metamorphism of Tibetan Ophiolitic Peridotites and its Tectonic Implications. *Journal of Petrology*, 57: 655-684. <https://doi.org/10.1093/petrology/egw011>.
- Grove T.L., Kinzler R.J. and Bryan W.B., 1992. Fractionation of mid-ocean ridge basalt (MORB). *Geophysical Monograph Series*, 71: 281-310.
- Guilmette C., Hébert R., Dostal J., Indares A., Ullrich T., Bédard É. and Wang C., 2012. Discovery of a dismembered metamorphic sole in the Saga ophiolitic mélange, South Tibet: Assessing an Early Cretaceous disruption of the Neo-Tethyan supra-subduction zone and consequences on basin closing. *Gondwana Research*, 22: 398-414. <https://doi.org/10.1016/j.gr.2011.10.012>.
- Guilmette C., Hébert R., Wang C. and Villeneuve M., 2009. Geochemistry and geochronology of the metamorphic sole underlying the Xigaze Ophiolite, Yarlung Zangbo Suture Zone, South Tibet. *Lithos*, 112: 149-162. <https://doi.org/10.1016/j.lithos.2009.05.027>.
- Hansen L.N., Cheadle M.J., John B.E., Swapp S.M., Dick H.J.B., Tucholke B.E. and Tivey M.A., 2013. Mylonitic deformation at the Kane oceanic core complex: Implications for the rheological behavior of oceanic detachment faults. *Geochemistry, Geophysics, Geosystems*, 14: 3085-3108. <https://doi.org/10.1002/ggge.20184>.
- Harvey J., Gannoun A., Burton K.W., Rogers N.W., Alard O. and Parkinson I.J., 2006. Ancient melt extraction from the oceanic upper mantle revealed by Re–Os isotopes in abyssal peridotites from the Mid-Atlantic ridge. *Earth and Planetary Science Letters*, 244: 606-621. <https://doi.org/10.1016/j.epsl.2006.02.031>.
- Hébert R., Bezard R., Guilmette C., Dostal J., Wang C.S. and Liu Z.F., 2012. The Indus–Yarlung Zangbo ophiolites from Nanga Parbat to Namche Barwa syntaxes, southern Tibet: First synthesis of petrology, geochemistry, and geochronology with incidences on geodynamic reconstructions of Neo-Tethys. *Gondwana Research*, 22: 377-397. <https://doi.org/10.1016/j.gr.2011.10.013>.
- Hu X.M., Garzanti E., Wang J.G., Huang W.T., An W. and Webb A., 2016. The timing of India-Asia collision onset – Facts, theories, controversies. *Earth-Science Reviews*, 160: 264-299. <https://doi.org/10.1016/j.earscirev.2016.07.014>.
- Huang W.T., Garzanti E., Moore T.C., Kapp P., DeCelles P.G. and Wang C.S., 2024. Continuity of the Sangdanlin Paleocene section and rejection of a large Greater India in the Early Cretaceous. *Proceedings of the National Academy of Sciences of the United States of America*, 121: e2402456121. <https://doi.org/10.1073/pnas.2402456121>.
- Ishizuka O., Tani K., Reagan M.K., Kanayama K., Umino S., Harigane Y., Sakamoto I., Miyajima Y., Yuasa M. and Dunkley D.J., 2011. The timescales of subduction initiation and subsequent evolution of an oceanic island arc. *Earth and Planetary Science Letters*, 306: 229-240. <https://doi.org/10.1016/j.epsl.2011.04.006>.
- Kempton P.D., Pearce J.A., Barry T.L., Fitton J.G., Langmuir C. and Christie D.M., 2002. Sr-Nd-Pb-Hf Isotope Results from ODP Leg 187: Evidence for Mantle Dynamics of the Australian-Antarctic Discordance and Origin of the Indian MORB Source. *Geochemistry, Geophysics, Geosystems*, 3: 1-35. <https://doi.org/10.1029/2002GC000320>.
- Kepezhinskas P. and Defant M. J., 2001. Nonchondritic Pt/Pd ratios in arc mantle xenoliths: Evidence for platinum enrichment in depleted island-arc mantle sources. *Geology*, 29: 851-854. [https://doi.org/10.1130/0091-7613\(2001\)029<0851:NPPRIA>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0851:NPPRIA>2.0.CO;2).
- Koepke J., Garbe-Schönberg D., Müller T., Mock D., Müller S. and Nasir S., 2022. A Reference Section Through Fast-Spread Lower Oceanic Crust, Wadi Gideah, Samail Ophiolite (Sultanate of Oman): Petrography and Petrology. *Journal of Geophysical Research: Solid Earth*, 127: e2021JB022735. <https://doi.org/10.1029/2021JB022735>.
- Lassiter J., Byerly B., Snow J. and Hellebrand E., 2014. Constraints from Os-isotope variations on the origin of Lena Trough abyssal peridotites and implications for the composition and evolution of the depleted upper mantle. *Earth and Planetary Science Letters*, 403: 178-187. <https://doi.org/10.1016/j.epsl.2014.05.033>.

- Li J., Jian H., Chen Y.J., Singh S.C., Ruan A., Qiu X., Zhao M., Wang X., Niu X., Ni J. and Zhang J., 2015. Seismic observation of an extremely magmatic accretion at the ultraslow spreading Southwest Indian Ridge. *Geophysical Research Letters*, 42: 2656-2663. <https://doi.org/10.1002/2014GL062521>.
- Li X.N., Zhang W.Q., Liu T., Zhang C., Lin Y.Z., Zhang Z., Zhu J.N., Li H.Y. and Liu C.Z., 2025a. Mantle evolution beneath a back-arc basin: Highly siderophile elements and Os isotope of Dongco ophiolitic peridotites, Tibet. *Chemical Geology*, 683: 122757. <https://doi.org/10.1016/j.chemgeo.2025.122757>.
- Li X.P., Zhang L., Wei C., Ai Y. and Chen J., 2007. Petrology of rodingite derived from eclogite in western Tianshan, China. *Journal of Metamorphic Geology*, 25: 363-382. <https://doi.org/10.1111/j.1525-1314.2007.00700.x>.
- Li Y., Li R.B., Dong T.C., Yang S.B. and Pei L., 2016. Structure of the Baimarang massif in the Xigaze ophiolite, Yarlung Zangbo Suture Zone, southern Tibet, China. *Chinese Science Bulletin*, 61: 2823-2833. <https://engine.scichina.com/doi/10.1360/N972016-00203>.
- Li Y., Li R.B., Liu C., Zhang J., Ciren D., Zhou X., He Z. and Yang J., 2025b. Detachment fault system in the Zedang ophiolite of southern Tibet: New insights into the evolution of the neo-Tethys ocean. *Journal of Asian Earth Sciences*, 283: 106525. <https://doi.org/10.1016/j.jseaes.2025.106525>.
- Li Y., Li R.B., Robinson P.T., Yang S.B., Zhang J. and Yang J.S., 2021. Detachment faulting in the Xigaze ophiolite southern Tibet: New constraints on its origin and implications. *Gondwana Research*, 94: 44-55. <https://doi.org/10.1016/j.gr.2021.02.006>.
- Lian D., Yang J., Robinson P.T., Liu F., Xiong F., Zhang L., Gao J. and Wu W., 2016. Tectonic Evolution of the Western Yarlung Zangbo Ophiolitic Belt, Tibet: Implications from the Petrology, Mineralogy, and Geochemistry of the Peridotites. *The Journal of Geology*, 124: 353-376.
- Lian D., Liu F., Cai P., Wu W., Li J., Majka J., Xu Z. and Yang J., 2024. Osmium and zinc isotope constraints on the origin of chromitites from the Yarlung-Zangbo ophiolites, Tibet, China. *Mineralium Deposita*, 59: 1089-1107. <https://doi.org/10.1007/s00126-024-01252-9>.
- Ligi M., Cuffaro M., Muccini F. and Bonatti E., 2022. Generation and evolution of the oceanic lithosphere in the North Atlantic. *La Rivista del Nuovo Cimento*, 45: 587-659. <https://doi.org/10.1007/s40766-022-00035-0>.
- Lin J. and Morgan J.P., 1992. The spreading rate dependence of three-dimensional mid-ocean ridge gravity structure. *Geophysical Research Letters*, 19: 13-16. <https://doi.org/10.1029/91GL03041>.
- Lin J., Purdy G.M., Schouten H., Sempere J.C. and Zervas C., 1990. Evidence from gravity data for focused magmatic accretion along the Mid-Atlantic Ridge. *Nature*, 344: 627-632. <https://doi.org/10.1038/344627a0>.
- Lin Y.Z., Day J.M., Brown D.B., Harvey J. and Liu C.Z., 2024. Evidence for large-scale, long-term highly siderophile element heterogeneities in the Atlantic mantle from Leg 153 and 209 peridotites. *Geochimica et Cosmochimica Acta*, 378: 300-314. <https://doi.org/10.1016/j.gca.2024.06.035>.
- Lissenberg C.J. and Dick H.J.B., 2008. Melt-rock reaction in the lower oceanic crust and its implications for the genesis of mid-ocean ridge basalt. *Earth and Planetary Science Letters*, 271: 311-325. <https://doi.org/10.1016/j.epsl.2008.04.023>.
- Lissenberg C.J. and MacLeod C.J., 2016. A Reactive Porous Flow Control on Mid-ocean Ridge Magmatic Evolution. *Journal of Petrology*, 57: 2195-2220. <https://doi.org/10.1093/petrology/egw074>.
- Lissenberg C.J., MacLeod C.J., Howard K.A. and Godard M., 2013. Pervasive reactive melt migration through fast-spreading lower oceanic crust (Hess Deep, equatorial Pacific Ocean). *Earth and Planetary Science Letters*, 361: 436-447. <https://doi.org/10.1016/j.epsl.2012.11.012>.
- Liu C.Z., Dick H.J., Mitchell R.N., Wei W., Zhang Z.Y., Hofmann A.W., Yang J.F. and Li Y., 2022a. Archean cratonic

- mantle recycled at a mid-ocean ridge. *Science Advances*, 8: eabn6749. <https://doi.org/10.1126/sciadv.abn6749>.
- Liu C.Z., Snow J.E., Brüggemann G., Hellebrand E. and Hofmann A.W., 2009. Non-chondritic HSE budget in Earth's upper mantle evidenced by abyssal peridotites from Gakkel ridge (Arctic Ocean). *Earth and Planetary Science Letters*, 283: 122-132. <https://doi.org/10.1016/j.epsl.2009.04.002>.
- Liu C.Z., Snow J.E., Hellebrand E., Brüggemann G., Von Der Handt A., Büchl A. and Hofmann A.W., 2008. Ancient, highly heterogeneous mantle beneath Gakkel ridge, Arctic Ocean. *Nature*, 452: 311-316. <https://doi.org/10.1038/nature06688>.
- Liu C.Z., Wu F.Y., Chu Z.Y., Ji W.Q., Yu L.J. and Li J.L., 2012. Preservation of ancient Os isotope signatures in the Yungbwa ophiolite (southwestern Tibet) after subduction modification. *Journal of Asian Earth Sciences*, 53: 38-50. <https://doi.org/10.1016/j.jseaes.2011.08.010>.
- Liu C.Z., Wu F.Y., Liu T., Zhang C., Zhang W.Q., Zhang Z.Y., Zhang Z., Wei W. and Lin Y.Z., 2022b. An origin of ultraslow spreading ridges for the Yarlung-Tsangpo ophiolites. *Fundamental Research*, 2: 74-83. <https://doi.org/10.1016/j.fmre.2021.07.002>.
- Liu C.Z., Zhang C., Yang L.Y., Zhang L.L., Ji W.Q. and Wu F.Y., 2014. Formation of gabbro-norites in the Purang ophiolite (SW Tibet) through melting of hydrothermally altered mantle along a detachment fault. *Lithos*, 205: 127-141. <https://doi.org/10.1016/j.lithos.2014.06.019>.
- Liu F., Dilek Y., Xie Y., Yang J. and Lian D., 2018a. Melt evolution of upper mantle peridotites and mafic dikes in the northern ophiolite belt of the western Yarlung Zangbo suture zone (southern Tibet). *Lithosphere*, 10: 109-132. <https://doi.org/10.1130/L689.1>.
- Liu T., Dick H.J.B., Liu C.Z., Wu F.Y., Ji W.B., Zhang C., Zhang W.Q., Zhang Z.Y., Lin Y.Z. and Zhang Z., 2021a. Tectonic Controls on Block Rotation and Sheeted Sill Emplacement in the Xigaze Ophiolite (Tibet): The Construction Mode of Slow-Spreading and Ultraslow-Spreading Oceanic Crusts. *Geochemistry, Geophysics, Geosystems*, 22: e2020GC009297. <https://doi.org/10.1029/2020GC009297>.
- Liu T., Liu C.Z., Wu F.Y., Dick H.J.B., Ji W.B., Zhang C., Zhang W.Q., Zhang Z.Y. and Xu Y., 2021b. The Xigaze ophiolite: fossil ultraslow-spreading ocean lithosphere in the Tibetan Plateau. *Journal of the Geological Society*, 178: jgs2020-2208. <https://doi.org/10.1144/jgs2020-208>.
- Liu T., Liu C.Z., Wu F.Y., Ji W.B., Zhang C., Zhang W.Q. and Zhang Z.Y., 2023. Timing and mechanism of opening the Neo-Tethys Ocean: Constraints from mélanges in the Yarlung Zangbo suture zone. *Science China Earth Sciences*, 66: 2807-2826. <https://doi.org/10.1007/s11430-023-1175-5>.
- Liu T., Liu C.Z., Wu F.Y., Topuz G., Wan B., Wang J.M. and Chen G., 2024. Detachment Fault-Hosted Subduction Re-Initiation of the (Ultra)Slow-Spreading Western Neo-Tethys in the Jurassic. *Geochemistry, Geophysics, Geosystems*, 25: e2023GC011173. <https://doi.org/10.1029/2023GC011173>.
- Liu T., Wu F.Y., Liu C.Z., Tribuzio R., Ji W.B., Zhang C., Xu Y. and Zhang W.Q., 2018b. Variably evolved gabbroic intrusions within the Xigaze ophiolite (Tibet): new insights into the origin of ophiolite diversity. *Contributions to Mineralogy and Petrology*, 173: 91. <https://doi.org/10.1007/s00410-018-1518-6>.
- Liu T., Wu F.Y., Liu C.Z., Zhang C., Ji W.B. and Xu Y., 2019. Reconsideration of Neo-Tethys evolution constrained from the nature of the Dazhuqu ophiolitic mantle, southern Tibet. *Contributions to Mineralogy and Petrology*, 174: 23. <https://doi.org/10.1007/s00410-019-1557-7>.
- Liu T., Wu F.Y., Liu C.Z., Zhu D.C. and Lin Y.Z., 2021c. Recycling of ancient sub-oceanic mantle in the Neo-Tethyan asthenosphere: Evidence from major and trace elements and Hf-Os isotopes of the Kop Mountain ophiolite, NE Turkey. *Geochimica et Cosmochimica Acta*, 311: 43-58. <https://doi.org/10.1016/j.gca.2021.08.003>.

- Liu T., Wu F.Y., Zhang L.L., Zhai Q.G., Liu C.Z., Ji W.B., Zhang C. and Xu Y., 2016. Zircon U-Pb geochronological constraints on rapid exhumation of the mantle peridotite of the Xigaze ophiolite, southern Tibet. *Chemical Geology*, 443: 67-86. <https://doi.org/10.1016/j.chemgeo.2016.09.015>.
- Lorand J.-P., Pattou L. and Gros M., 1999. Fractionation of Platinum-group Elements and Gold in the Upper Mantle: a Detailed Study in Pyrenean Orogenic Lherzolites. *Journal of Petrology*, 40: 957-981. <https://doi.org/10.1093/petroj/40.6.957>.
- Luguet A. and Reisberg L., 2016. Highly Siderophile Element and ^{187}Os Signatures in Non-cratonic Basalt-hosted Peridotite Xenoliths: Unravelling the Origin and Evolution of the Post-Archean Lithospheric Mantle. *Reviews in Mineralogy and Geochemistry*, 81: 305-367.
- MacLeod C.J., Carlot J., Escartín J., Horen H. and Morris A., 2011. Quantitative constraint on footwall rotations at the 15° 45'N oceanic core complex, Mid-Atlantic Ridge: Implications for oceanic detachment fault processes. *Geochemistry, Geophysics, Geosystems*, 12: Q0AG03. <https://doi.org/10.1029/2011GC003503>.
- MacLeod C.J., Escartín J., Banerji D., Banks G.J., Gleeson M., Irving D.H.B., Lilly R.M., McCaig A.M., Niu Y., Allerton S. and Smith D.K., 2002. Direct geological evidence for oceanic detachment faulting: The Mid-Atlantic Ridge, 15° 45'N. *Geology*, 30: 879-882. [https://doi.org/10.1130/0091-7613\(2002\)030<0879:DGEFOD>2.0.CO;2](https://doi.org/10.1130/0091-7613(2002)030<0879:DGEFOD>2.0.CO;2).
- MacLeod C.J. and Yaouancq G., 2000. A fossil melt lens in the Oman ophiolite: Implications for magma chamber processes at fast spreading ridges. *Earth and Planetary Science Letters*, 176: 357-373. [https://doi.org/10.1016/S0012-821X\(00\)00020-0](https://doi.org/10.1016/S0012-821X(00)00020-0).
- Maffione M., Morris A. and Anderson M.W., 2013. Recognizing detachment-mode seafloor spreading in the deep geological past. *Scientific reports*, 3: 2336. <https://doi.org/10.1038/srep02336>.
- Maffione M., van Hinsbergen D.J.J., Koornneef L.M.T., Guilmette C., Hodges K., Borneman N., Huang W., Ding L. and Kapp P., 2015. Forearc hyperextension dismembered the south Tibetan ophiolites. *Geology*, 43: 475-478. <https://doi.org/10.1130/G36472.1>.
- Mahoney J., le Roex A.P., Peng Z., Fisher R.L. and Natland J.H., 1992. Southwestern limits of Indian Ocean Ridge Mantle and the origin of low $^{206}\text{Pb}/^{204}\text{Pb}$ mid-ocean ridge basalt: Isotope systematics of the central Southwest Indian Ridge (17°–50°E). *Journal of Geophysical Research: Solid Earth*, 97: 19771-19790. <https://doi.org/10.1029/92JB01424>.
- Malpas J., Zhou M.F., Robinson P.T. and Reynolds P.H., 2003. Geochemical and geochronological constraints on the origin and emplacement of the Yarlung Zangbo ophiolites, Southern Tibet. In: Y. Dilek and P.T. Robinson (Eds.), *Ophiolites in Earth History*. Geological Society of London, 218: 191-206. <https://doi.org/10.1144/GSL.SP.2003.218.01.11>.
- Manatschal G., Sauter D., Karpoff A.M., Masini E., Mohn G. and Lagabrielle Y., 2011. The Chenaillet Ophiolite in the French/Italian Alps: An ancient analogue for an Oceanic Core Complex? *Lithos*, 124, 169-184. <https://doi.org/10.1016/j.lithos.2010.10.017>.
- McCarthy A. and Münster O., 2015. Ancient depletion and mantle heterogeneity: Revisiting the Permian-Jurassic paradox of Alpine peridotites. *Geology*, 43: 255-258. <https://doi.org/10.1130/G36340.1>.
- McKenzie D. and O'Nions R.K., 1983. Mantle reservoirs and ocean island basalts. *Nature*, 301: 229-231. <https://doi.org/10.1038/301229a0>.
- Meisel T., Walker R.J., Irving A.J. and Lorand J.-P., 2001. Osmium isotopic compositions of mantle xenoliths: a global perspective. *Geochimica et Cosmochimica Acta*, 65: 1311-1323. [https://doi.org/10.1016/S0016-7037\(00\)00566-4](https://doi.org/10.1016/S0016-7037(00)00566-4).
- Mendel V., Sauter D., Parson L. and Vanney J.-R., 1997. Segmentation and Morphotectonic Variations Along a Super Slow-Spreading Center: The Southwest Indian Ridge (57° E-70° E). *Marine Geophysical Researches*, 19: 505-533.

<https://doi.org/10.1023/A:1004232506333>.

- Miyashiro A., 1973. The Troodos ophiolitic complex was probably formed in an island arc. *Earth and Planetary Science Letters*, 19: 218-224. [https://doi.org/10.1016/0012-821X\(73\)90118-0](https://doi.org/10.1016/0012-821X(73)90118-0).
- Moore E.M., Kellogg L.H. and Dilek, Y., 2000. Tethyan ophiolites, mantle convection, and tectonic “historical contingency”: A resolution of the “ophiolite conundrum”. In: Y. Dilek, E.M. Moore, D. Elthon and A. Nicolas (Eds.), *Ophiolites and Oceanic Crust: New Insights from Field Studies and the Ocean Drilling Program*. Boulder, Colorado, Geological Society of America Special Paper 349: 3–12. <https://doi.org/10.1130/0-8137-2349-3.3>.
- Moore E.M., Simmons N., Basu A.R. and Gregory R.T., 2021. The Indian Ocean, its supra-subduction history, and implications for ophiolites. In: J. Wakabayashi and Y. Dilek (Eds.), *Plate Tectonics, Ophiolites, and Societal Significance of Geology: A Celebration of the Career of Eldridge Moore*. Geological Society of America Special Paper, 552: 1–14. [https://doi.org/10.1130/2021.2552\(01\)](https://doi.org/10.1130/2021.2552(01)).
- Morris A., Gee J.S., Pressling N., John B.E., MacLeod C.J., Grimes C.B. and Searle R.C., 2009. Footwall rotation in an oceanic core complex quantified using reoriented Integrated Ocean Drilling Program core samples. *Earth and Planetary Science Letters*, 287: 217-228. <https://doi.org/10.1016/j.epsl.2009.08.007>.
- Müller S., Garbe-Schönberg D., Koepke J. and Hoernle K., 2022. A Reference Section Through Fast-Spread Lower Oceanic Crust, Wadi Gideah, Samail Ophiolite (Sultanate of Oman): Trace Element Systematics and REE Crystallization Temperatures—Implications for Hybrid Crustal Accretion. *Journal of Geophysical Research: Solid Earth*, 127: e2021JB022699. <https://doi.org/10.1029/2021JB022699>.
- Müntener O. and Manatschal G., 2006. High degrees of melt extraction recorded by spinel harzburgite of the Newfoundland margin: the role of inheritance and consequences for the evolution of the southern North Atlantic. *Earth and Planetary Science Letters*, 252: 437-452. <https://doi.org/10.1016/j.epsl.2006.10.009>.
- Nauret F., Snow J.E., Hellebrand E. and Weis D., 2011. Geochemical Composition of K-rich Lavas from the Lena Trough (Arctic Ocean). *Journal of Petrology*, 52: 1185-1206. <https://doi.org/10.1093/petrology/egr024>.
- Nicolas A., Boudier F. and Meshi A., 1999. Slow spreading accretion and mantle denudation in the Mirdita ophiolite (Albania). *Journal of Geophysical Research*, 104: 15155-15167. <https://doi.org/10.1029/1999JB900126>.
- Nicolas A., Girardeau J., Marcoux J., Dupre B., Xubin W., Yougong C., Haixiang Z. and Xuchang X., 1981. The Xigaze ophiolite (Tibet): a peculiar oceanic lithosphere. *Nature*, 294: 414-417. <https://doi.org/10.1038/294414a0>.
- Niu X., Yang J., Dilek Y., Xu J., Li J., Chen S., Feng G., Liu F., Xiong F. and Liu Z., 2015. Petrological and Os isotopic constraints on the origin of the Dongbo peridotite massif, Yarlung Zangbo Suture Zone, Western Tibet. *Journal of Asian Earth Sciences*, 110: 72-84. <https://doi.org/10.1016/j.jseas.2014.09.036>.
- Ou G.F., Wang S.J., Deng H., Duan W.Y., Li X.P., Schertl H.-P., 2025. Fore-arc metasomatism by hybrid slab fluids during subduction initiation: Sr-Mg-Ca isotopes of rodingite, western Yarlung Zangbo suture zone. *Chemical Geology*, 671: 122473. <https://doi.org/10.1016/j.chemgeo.2024.122473>.
- Parlak O., Çolakoğlu A., Dönmez C., Sayak H., Yildirim N., Türkel A. and Odabaşı İ., 2012. Geochemistry and tectonic significance of ophiolites along the İzmir-Ankara-Erzincan Suture Zone in northeastern Anatolia. Geological Society, London, Special Publications, 372, <https://doi.org/10.1144/SP372.7>.
- Parlak O., Karaoğlu F., Rızaoğlu T., Klötzli U., Koller F., and Billor Z., 2013. U–Pb and ⁴⁰Ar–³⁹Ar geochronology of the ophiolites and granitoids from the Tauride belt: Implications for the evolution of the Inner Tauride suture. *Journal of Geodynamics*, 65: 22-37. <https://doi.org/10.1016/j.jog.2012.06.012>.
- Pearce J.A., 2014. Immobile Element Fingerprinting of Ophiolites. *Elements*, 10: 101-108. <https://doi.org/10.2113/gselements.10.2.101>.

- Rampone E. and Hofmann A.W., 2012. A global overview of isotopic heterogeneities in the oceanic mantle. *Lithos*, 148: 247-261. <https://doi.org/10.1016/j.lithos.2012.06.018>.
- Rampone E. and Sanfilippo A., 2021. The Heterogeneous Tethyan Oceanic Lithosphere of the Alpine Ophiolites. *Elements*, 17, 23-28. <https://doi.org/10.2138/gselements.17.1.23>.
- Reagan M.K., Ishizuka O., Stern R.J., Kelley K.A., Ohara Y., Blichert-Toft J., Bloomer S.H., Cash J., Fryer P., Hanan B.B., Hickey-Vargas R., Ishii T., Kimura J.-I., Peate D.W., Rowe M.C. and Woods M., 2010. Fore-arc basalts and subduction initiation in the Izu-Bonin-Mariana system. *Geochemistry, Geophysics, Geosystems*, 11: Q03X12. <https://doi.org/10.1029/2009GC002871>.
- Rommevaux-Jestin C., Deplus C. and Patriat P., 1997. Mantle Bouguer Anomaly Along an Ultra Slow-Spreading Ridge: Implications for Accretionary Processes and Comparison with Results from Central Mid-Atlantic Ridge. *Marine Geophysical Researches*, 19: 481-503. <https://doi.org/10.1023/A:1004269003009>.
- Rudnick R.L. and Walker R.J., 2009. Interpreting ages from Re–Os isotopes in peridotites. *Lithos*, 112: 1083-1095. <https://doi.org/10.1016/j.lithos.2009.04.042>.
- Sanfilippo A., Dick H.J.B., Marschall H.R., Lissenberg C.J. and Urann B., 2018. Emplacement and High-Temperature Evolution of Gabbros of the 16.5°N Oceanic Core Complexes (Mid-Atlantic Ridge): Insights Into the Compositional Variability of the Lower Oceanic Crust. *Geochemistry, Geophysics, Geosystems*, 20: 2018GC007512. <https://doi.org/10.1029/2018GC007512>.
- Sanfilippo A., MacLeod C.J., Tribuzio R., Lissenberg C.J. and Zanetti A., 2020. Early-Stage Melt-Rock Reaction in a Cooling Crystal Mush Beneath a Slow-Spreading Mid-Ocean Ridge (IODP Hole U1473A, Atlantis Bank, Southwest Indian Ridge). *Frontiers in Earth Science*, 8: 579138. <https://doi.org/10.3389/feart.2020.579138>.
- Sanfilippo A., Stracke A., Genske F., Scarani S., Cuffaro M., Basch V., Borghini G., Brunelli D., Ferrando C., Peyve A.A. and Ligi M., 2024. Upwelling of melt-depleted mantle under Iceland. *Nature Geoscience*, 17: 1046-1052. <https://doi.org/10.1038/s41561-024-01532-z>.
- Sanfilippo A. and Tribuzio R., 2011. Melt transport and deformation history in a nonvolcanic ophiolitic section, northern Apennines, Italy: Implications for crustal accretion at slow spreading settings. *Geochemistry, Geophysics, Geosystems*, 12: Q0AG04. <https://doi.org/10.1029/2010GC003429>.
- Sanfilippo A. and Tribuzio R., 2013. Building of the deepest crust at a fossil slow-spreading centre (Pineto gabbroic sequence, Alpine Jurassic ophiolites). *Contributions to Mineralogy and Petrology*, 165: 705-721. <https://doi.org/10.1007/s00410-012-0831-8>.
- Sani C., Sanfilippo A., Peyve A.A., Genske F. and Stracke A., 2023. Earth Mantle's Isotopic Record of Progressive Chemical Depletion. *AGU Advances*, 4: e2022AV000792. <https://doi.org/10.1029/2022AV000792>.
- Sarifakioğlu E., Özen H. and Winchester J.A., 2009. Petrogenesis of the Refahiye ophiolite and its Tectonic significance for Neotethyan Ophiolites Along the İzmir-Ankara-Erzincan Suture Zone. *Turkish Journal of Earth Sciences*, 18: 187-207. <https://doi.org/10.3906/yer-0702-4>.
- Sinton J.M. and Detrick R.S., 1992. Mid-ocean ridge magma chambers. *Journal of Geophysical Research: Solid Earth*, 97: 197-216. <https://doi.org/10.1029/91JB02508>.
- Stern R.J., 2004. Subduction initiation: spontaneous and induced. *Earth and Planetary Science Letters*, 226: 275-292. <https://doi.org/10.1016/j.epsl.2004.08.007>.
- Stern R. J., Reagan M., Ishizuka O., Ohara Y. and Whattam S., 2012. To understand subduction initiation, study forearc crust: To understand forearc crust, study ophiolites. *Lithosphere*, 4: 469-483. <https://doi.org/10.1130/L183.1>.
- Stracke A., 2021. A process-oriented approach to mantle geochemistry. *Chemical Geology*, 579: 120350.

- <https://doi.org/10.1016/j.chemgeo.2021.120350>.
- Stracke A., Genske F., Berndt J. and Koornneef J.M., 2019. Ubiquitous ultra-depleted domains in Earth's mantle. *Nature Geoscience*, 12: 851-855. <https://doi.org/10.1038/s41561-019-0446-z>.
- Stracke A., Snow J.E., Hellebrand E., von der Handt A., Bourdon B., Birbaum K. and Günther D., 2011. Abyssal peridotite Hf isotopes identify extreme mantle depletion. *Earth and Planetary Science Letters*, 308: 359-368. <https://doi.org/10.1016/j.epsl.2011.06.012>.
- Sun S.S. and McDonough W.F., 1989. Chemical and isotopic systematics of oceanic basalts: implications for mantle composition and processes. Geological Society, London, Special Publications, 42: 313-345. <https://doi.org/10.1144/GSL.SP.1989.042.01.19>.
- Tolstoy M., Harding A.J. and Orcutt J.A., 1993. Crustal Thickness on the Mid-Atlantic Ridge: Bull's-Eye Gravity Anomalies and Focused Accretion. *Science*, 262: 726-729. <https://doi.org/10.1126/science.262.5134.726>.
- Tucholke B.E., Behn M.D., Buck W.R. and Lin J., 2008. Role of melt supply in oceanic detachment faulting and formation of megamullions. *Geology*, 36: 455-458. <https://doi.org/10.1130/G24639A.1>.
- Urann B.M., Dick H.J.B., Parnell-Turner R. and Casey J.F., 2020. Recycled arc mantle recovered from the Mid-Atlantic Ridge. *Nature Communications*, 11: 3887. <https://doi.org/10.1038/s41467-020-17604-8>.
- Wakabayashi J. and Dilek, Y., 2003. What constitutes 'emplacement' of an ophiolite?: Mechanisms and relationship to subduction initiation and formation of metamorphic soles. Geological Society, London, Special Publications, 218: 427-447. <https://doi.org/10.1144/GSL.SP.2003.218.01.22>.
- Wang J., Zhou H., Salters V.J.M., Dick H.J.B., Standish J.J. and Wang C., 2020. Trace Element and Isotopic Evidence for Recycled Lithosphere from Basalts from 48 to 53°E, Southwest Indian Ridge. *Journal of Petrology*, 61: ega068. <https://doi.org/10.1093/petrology/egaa068>.
- Wang J.G., Hu X., Garzanti E., An W. and Liu X.C., 2017. The birth of the Xigaze forearc basin in southern Tibet. *Earth and Planetary Science Letters*, 465: 38-47. <https://doi.org/10.1016/j.epsl.2017.02.036>.
- Wang X.B., Bao P.S., Deng W.M. and Wang F.G., 1987. Xigaze (Tibet) Ophiolites. Beijing, Geological Publishing House: 1-336 (in Chinese).
- Wang Y., Mungall J.E. and Liu J., 2023. Osmium isotope heterogeneity of the upper mantle: Evidence from the Bay of Islands ophiolite complex, Newfoundland. *Journal of Geophysical Research: Solid Earth*, 128: e2023JB026789. <https://doi.org/10.1029/2023JB026789>.
- Warren J.M., 2016. Global variations in abyssal peridotite compositions. *Lithos*, 248-251: 193-219. <https://doi.org/10.1016/j.lithos.2015.12.023>.
- Whattam S.A. and Stern R.J., 2011. The 'subduction initiation rule': a key for linking ophiolites, intra-oceanic forearcs, and subduction initiation. *Contributions to Mineralogy and Petrology*, 162: 1031-1045. <https://doi.org/10.1007/s00410-011-0638-z>.
- Whitehead J.A., Dick H.J.B. and Schouten H., 1984. A mechanism for magmatic accretion under spreading centres. *Nature*, 312: 146-148. <https://doi.org/10.1038/312146a0>.
- Wu F.Y., Liu C.Z., Zhang L.L., Zhang C., Wang J.G., Ji W.Q. and Liu X.C., 2014. Yanglung Zangbo ophiolite: A critical updated view. *Acta Petrologica Sinica*, 30: 293-325 (in Chinese with English abstract).
- Wu F.Y., Wan B., Zhao L., Xiao W.J. and Zhu R.X., 2020. Tethyan geodynamics. *Acta Petrologica Sinica*, 36: 1627-1674 (in Chinese with English abstract).
- Xia B., Yu H.X., Chen G.W., Qi L., Zhao T.P. and Zhou M.F., 2003. Geochemistry and tectonic environment of the Dagzhuka ophiolite in the Yarlung-Zangbo suture zone, Tibet. *Geochemical Journal*, 37: 311-324.

<https://doi.org/10.2343/geochemj.37.311>.

- Xin G.Y., Chu Y., Su B.X., Lin W., Uysal I. and Feng Z.T., 2021. Rapid transition from MORB-type to SSZ-type oceanic crust generation following subduction initiation: insights from the mafic dikes and metamorphic soles in the Pozanti-Karsanti ophiolite, SE Turkey. *Contributions to Mineralogy and Petrology*, 176: 64. <https://doi.org/10.1007/s00410-021-01821-5>.
- Xiong F., Yang J., Robinson P.T., Gao J., Chen Y. and Lai S., 2017a. Petrology and geochemistry of peridotites and podiform chromitite in the Xigaze ophiolite, Tibet: Implications for a suprasubduction zone origin. *Journal of Asian Earth Sciences*, 146: 56-75. <https://doi.org/10.1016/j.jseaes.2017.05.001>.
- Xiong Q., Dai H.K., Zheng J.P., Griffin W.L., Zheng H.D., Wang L. and Reilly S.Y.O., 2022. Vertical depletion of ophiolitic mantle reflects melt focusing and interaction in sub-spreading-center asthenosphere. *Nature Communications*, 13: 6956. <https://doi.org/10.1038/s41467-022-34781-w>.
- Xiong Q., Griffin W.L., Zheng J.P., Pearson N.J. and O'Reilly S.Y., 2017b. Two-layered oceanic lithospheric mantle in a Tibetan ophiolite produced by episodic subduction of Tethyan slabs. *Geochemistry, Geophysics, Geosystems*, 18: 1189-1213. <https://doi.org/10.1002/2016GC006681>.
- Xiong Q., Griffin W.L., Zheng J.P., O'Reilly S.Y., Pearson N.J., Xu B. and Belousova E.A., 2016. Southward trench migration at ~130–120 Ma caused accretion of the Neo-Tethyan forearc lithosphere in Tibetan ophiolites. *Earth and Planetary Science Letters*, 438: 57-65. <https://doi.org/10.1016/j.epsl.2016.01.014>.
- Xu Y. and Liu C.Z., 2019. Subduction-Induced Fractionated Highly Siderophile Element Patterns in Forearc Mantle. *Minerals*, 9: 339. <https://doi.org/10.3390/min9060339>.
- Xu Y., Liu C.Z., Zhang C. and Liu T., 2025. Re-Os isotopic evidence for ancient melt depletion in refertilized Neo-Tethyan suboceanic mantle domain. *Chemical Geology*, 673: 122520. <https://doi.org/10.1016/j.chemgeo.2024.122520>.
- Xu Y., Liu J., Xiong Q., Su B.X., Scott J.M., Xu B., Zhu D.C. and Pearson D.G., 2020. The complex life cycle of oceanic lithosphere: A study of Yarlung-Zangbo ophiolitic peridotites, Tibet. *Geochimica et Cosmochimica Acta*, 277: 175-191. <https://doi.org/10.1016/j.gca.2020.03.024>.
- Yang A.Y., Langmuir C.H., Cai Y., Michael P., Goldstein S.L. and Chen Z., 2021. A subduction influence on ocean ridge basalts outside the Pacific subduction shield. *Nature Communications*, 12: 4757. <https://doi.org/10.1038/s41467-021-25027-2>.
- Yang S., Humayun M. and Salters V.J.M., 2020. Elemental constraints on the amount of recycled crust in the generation of mid-oceanic ridge basalts (MORBs). *Science Advances*, 6: eaba2923. <https://doi.org/10.1126/sciadv.aba2923>.
- Yin A. and Harrison T.M., 2000. Geologic Evolution of the Himalayan-Tibetan Orogen. *Annual Review of Earth and Planetary Sciences*, 28: 211-280. <https://doi.org/10.1146/annurev.earth.28.1.211>.
- Zhang C., Liu C.Z., Bénard A., Müntener O., Ji W.B., Liu T., Zhang Z.Y., Zhang W.Q. and Wu F.Y., 2023a. Heterogeneous mantle beneath the Neo-Tethys Ocean revealed by ultramafic rocks from the Xiugugabu Ophiolite in the Yarlung-Tsangpo Suture Zone, southwestern Tibet. *Contributions to Mineralogy and Petrology*, 178: 54. <https://doi.org/10.1007/s00410-023-02039-3>.
- Zhang C., Liu C.Z., Ji W.B., Liu T. and Wu F.Y., 2020a. Heterogeneous sub-ridge mantle of the Neo-Tethys: Constraints from Re-Os isotope and HSE compositions of the Xigaze ophiolites. *Lithos*, 378-379: 105819. <https://doi.org/10.1016/j.lithos.2020.105819>.
- Zhang C., Liu C.Z., Ji W.B., Liu T., Zhang Z.Y. and Wu F.Y., 2023b. Subduction of solidifying melt lens beneath ultraslow-spreading ridges revealed by (Grt-)amphibolites and dolerites in the Xiugugabu ophiolitic mélange, southwestern Tibet. *Lithos*, 462-463: 107421. <https://doi.org/10.1016/j.lithos.2023.107421>.

- Zhang C., Liu C.Z., Liu T., Ji W.B. and Wu F.Y., 2023c. Generation of ultraslow-spreading oceanic crust traced by various mafic blocks from ophiolitic mélange in the Xigaze Ophiolites, southern Tibet. *Contributions to Mineralogy and Petrology*, 178: 55. <https://doi.org/10.1007/s00410-023-02040-w>.
- Zhang C., Liu C.Z., Liu T. and Wu F.Y., 2020b. Evolution of mantle peridotites from the Luobusa ophiolite in the Tibetan Plateau: Sr-Nd-Hf-Os isotope constraints. *Lithos*, 362-363: 105477. <https://doi.org/10.1016/j.lithos.2020.105477>.
- Zhang C., Liu C.Z., Wu F.Y., Ji W.B., Liu T. and Xu Y., 2017. Ultra-refractory mantle domains in the Luqu ophiolite (Tibet): Petrology and tectonic setting. *Lithos*, 286-287: 252-263. <https://doi.org/10.1016/j.lithos.2017.05.021>.
- Zhang C., Liu C.Z., Wu F.Y., Zhang L.L. and Ji W.Q., 2016a. Geochemistry and geochronology of mafic rocks from the Luobusa ophiolite, South Tibet. *Lithos*, 245: 93-108. <https://doi.org/10.1016/j.lithos.2015.06.031>.
- Zhang C., Liu C.Z., Xu Y., Ji W.B., Wang J.M., Wu F.Y., Liu T., Zhang Z.Y. and Zhang W.Q., 2019. Subduction re-initiation at dying ridge of Neo-Tethys: Insights from mafic and metamafic rocks in Lhaze ophiolitic mélange, Yarlung-Tsangbo Suture Zone. *Earth and Planetary Science Letters*, 523: 115707. <https://doi.org/10.1016/j.epsl.2019.07.009>.
- Zhang L.L., Liu C.Z., Wu F.Y., Ji W.Q. and Wang J.G., 2014. Zedong terrane revisited: An intra-oceanic arc within Neo-Tethys or a part of the Asian active continental margin? *Journal of Asian Earth Sciences*, 80: 34-55. <https://doi.org/10.1016/j.jseaes.2013.10.029>.
- Zhang L.L., Liu C.Z., Wu F.Y., Zhang C., Ji W.Q. and Wang J.G., 2016b. Sr-Nd-Hf isotopes of the intrusive rocks in the Cretaceous Xigaze ophiolite, southern Tibet: Constraints on its formation setting. *Lithos*, 258-259: 133-148. <https://doi.org/10.1016/j.lithos.2016.04.026>.
- Zhang T., Li J., Niu X., Ding W., Fang Y., Lin J., Wang Y., Zha C., Tan P., Kong F., Chen J., Wei X., Lu J., Dymant J. and Morgan J.P., 2024a. Highly variable magmatic accretion at the ultraslow-spreading Gakkell Ridge. *Nature*, 633: 109-113. <https://doi.org/10.1038/s41586-024-07831-0>.
- Zhang W.Q., Dick H.J.B., Liu C.Z., Lin Y.Z. and Angeloni L.M., 2021. MORB Melt Transport through Atlantis Bank Oceanic Batholith (SW Indian Ridge). *Journal of Petrology*, 62: egab034. <https://doi.org/10.1093/petrology/egab034>.
- Zhang W.Q., Liu C.Z. and Dick H.J.B., 2020c. Evidence for Multi-stage Melt Transport in the Lower Ocean Crust: the Atlantis Bank Gabbroic Massif (IODP Hole U1473A, SW Indian Ridge). *Journal of Petrology*, 61: egaa082. <https://doi.org/10.1093/petrology/egaa082>.
- Zhang W.Q., Liu C.Z. and Liu T., 2022. Compositions of cumulates in lower oceanic crust and implications for ophiolite studies. *Acta Petrologica Sinica*, 38: 1630-1654 (in Chinese with English abstract).
- Zhang Z., Liu C.Z., Jung H., Liu T., Zhang C., Cao Y., Jung S., Zhang W.Q., Lin Y.Z. and Li X.N., 2025. Multi-Stage Evolution of Deformed Peridotites in the Yarlung-Tsangpo Suture Zone: A Link to Mantle Shear Initiation at H₂O-Deficient Ultraslow-Spreading Ridges. *Journal of Petrology*, 66: egaf073. <https://doi.org/10.1093/petrology/egaf073>.
- Zhang Z., Liu T., Liu C.Z., Jung S. and Jung H., 2023d. The water-poor and reduced mantle beneath the Neo-Tethys Ocean: Tectonic setting of the western Yarlung-Tsangpo ophiolites. *Lithos*, 456-457: 107326. <https://doi.org/10.1016/j.lithos.2023.107326>.
- Zhang Z.Y., Liu C.Z., Liang Y., Zhang L.L., Zhang C., Liu T., Liu B.D., Zhang W.Q., Lin Y.Z. and Ji W.B., 2024b. Petrogenesis of the pyroxenitic Moho transition zone in the Zedong ophiolite, Southeastern Tibet. *Lithos*, 482-483: 107703. <https://doi.org/10.1016/j.lithos.2024.107703>.
- Zhou B., Liu J., Yan J., Hou C., Chen X., Liu C. and Wu F., 2023. Shearing-enhanced deep fluid circulation induces seismic anisotropy in the lower crust at slow-spreading oceanic ridges. *Geology*, 51: 465-470. <https://doi.org/10.1130/G50879.1>.

- Zhou H. and Dick H.J.B., 2013. Thin crust as evidence for depleted mantle supporting the Marion Rise. *Nature*, 494: 195-200. <https://doi.org/10.1038/nature11842>.
- Zhu D.C., Zhao Z.D., Niu Y., Dilek Y., Hou Z.Q. and Mo X.X., 2013. The origin and pre-Cenozoic evolution of the Tibetan Plateau. *Gondwana Research*, 23: 1429-1454. <https://doi.org/10.1016/j.gr.2012.02.002>.
- Zhu J.N., Liu C.Z., Zhang W.Q., Liu T., Zhang C., Li X.N. and Zhang Z.Y., 2025. Recycling ancient refractory peridotites causing the crust-mantle decoupling of the Xigaze ophiolite (South Tibet): New constraints from the Buma mantle massif. *Lithos*, 508-509: 108090. <https://doi.org/10.1016/j.lithos.2025.108090>.
- Ziabrev S., Aitchison J., Abrajevitch A., Badengzhu, Davis A. and Luo H., 2003. Precise radiolarian age constraints on the timing of ophiolite generation and sedimentation in the Dazhuqu terrane, Yarlung–Tsangpo suture zone, Tibet. *Journal of the Geological Society*, 160: 591-599. <https://doi.org/10.1144/0016-764902-107>.

Pre-proof




Fig. 1. Geological framework of the Yarlung-Tsangpo ophiolites (YTO). (a) Tectonic framework of the Tibetan Plateau, showing the distribution and extension of major Tethyan sutures. AKMS: A'nyemaqen-Kunlun-Muztagh; JS: Jinsha suture; LSS: Longmuco-Shuanghu suture; BNS: Bangong-Nujiang suture; YTS: Yarlung-Tsangpo suture. (b) Tectonic units identified in southern Tibet and the distribution and segmentation of the YTO. (c) Zircon U-Pb dating results for different lithologies of the YTO and their comparisons with the ages of metamorphic soles and radiolarian cherts along the YTS. The age data is compiled in Table S1.




Fig. 2. Basic geochemical characteristics of the basaltic rocks (basalt and diabase; a-b), the lower crustal cumulates (c-d), and the mantle peridotites (e-g) of the YTO. Data for basalts, diabbases, and lower crustal cumulates is for the well-studied Xigaze ophiolite in the central segment of the YTO. Data for the mantle peridotites is from the entire suture. Data for N-MORB is from Sun and McDonough (1989). Average compositions of forearc basalts are after Reagan et al. (2010). Fields of oceanic and continental arcs are after Pearce (2014). The compositions of abyssal and SSZ cumulates are after the compilation by Zhang et al. (2022). Data for abyssal and forearc peridotites are from the PetDB database (<https://search.earthchem.org/>). N-MORB: normal mid-ocean ridge basalt; E-MORB: enriched mid-ocean ridge basalt; OIB: ocean island basalt; Pl: plagioclase; Cpx: clinopyroxene; Ol: olivine; Cr#: $Cr / (Cr + Al)$; Fo: $100 \times Mg / (Mg + Fe^{2+})$.




Fig. 3. Pseudostratigraphy of representative ophiolites along the Neo-Tethyan belt from the Alps to Tibet. References: Alps (Manatschal et al., 2011; Sanfilippo and Tribuzio, 2011, 2013), Mirdita (Dilek and Furnes, 2009; Nicolas et al., 1999), Vourinous (Bortolotti et al., 2004), Refahiye (Liu et al., 2024; Parlak et al., 2012; Sarifakioğlu et al., 2009), Inner Tauride (Parlak et al., 2013; Xin et al., 2021), Troodos & Kızıldağ & Baër-Bassit (Dilek and Thy, 2009), Oman (Dilek and Furnes, 2009), Kermanshah (Ao et al., 2016), Xigaze (Liu et al., 2021a, b; Nicolas et al., 1981), Purang (Liu et al., 2014).




Fig. 4. Field photos of the primary contacts at the bottom and on the top of the oceanic crust of the Xigaze ophiolite, central YTO. (a) Intrusive contact between lower crustal gabbros and serpentinized mantle peridotites (Dazhuqu). (b) Pillow lavas are directly overlain by radiolarian cherts (Naxia).



Fig. 5. Field relationships for the isolated mafic dikes/sills and sheeted sill complex in the Xigaze ophiolite, central YTO. (a) Serpentinized mantle peridotites are intruded by a fresh diabase dike, which displays clear two-way chilled margins (Pengcang). (b) Serpentinized mantle peridotites crosscut by several rodingite dikes (Ngamring). (c) Foliated amphibolites identified in the serpentinized mantle peridotites (Luqu). (d-e) Diabase dikes and sills intruding the lower crustal layered gabbros (Jiding). (f) The sheeted sill complex composed entirely of diabase sills (Jiding).




Fig. 6. (a) Geological map of the Xigaze ophiolite (modified after Wang et al., 1987). (b) Reconstructed oceanic crustal thicknesses along the paleo-ridge axis at the time of the ophiolite formation (~126 Ma; after Liu et al., 2021).




Fig. 7. (a) The measured ~350-m-thick lower crust in the Jiding ophiolite section, Xigaze ophiolite. (b) Schematic cryptic chemical variations of the Jiding lower crust (representing a low magma budget) and its comparison with the Atlantis Bank gabbro massif of the Southwest Indian Ridge (after Dick et al., 2002) and the Oman ophiolite (after Koepke et al. (2022) and MacLeod and Yaouancq (2000)). (c-f) Cryptic chemical variations of the Jiding gabbroic rocks.




Fig. 8. Along axis magma focusing (a) and cross-axis detachment faulting (b) identified for the paleo-ridge of the YTO, after Liu et al. (2016, 2021b).




Fig. 9. Reconstructed geochemical variations of the mantle peridotites along the ancient, >1300-km-long Neo-Tethyan mid-ocean ridge represented by the YTO. Melting extent is calculated by the empirical formula established based on spinel Cr# values of mantle peridotites (Warren, 2016).





Fig. 10. Re-Os isotopes and highly siderophile element (HSE) compositions for available data of the mantle peridotites within the YTO. Chondrite values for the normalization of HSE is after Fischer-Gödde et al. (2010). The composition of primitive upper mantle (PUM) is after Becker et al. (2006) and Meisel et al. (2001). Data for abyssal peridotites and forearc peridotites is from the PetDB database (<https://search.earthchem.org/>).

(a) Ultraslow-spreading ridge (~130-120 Ma) with a heterogeneous asthenosphere



(b) Subduction jump and re-initiation (<120 Ma) along detachment faults




Fig. 11. Two new conceptual models for alternative explanations of the YTO ophiolite conundrum. (a) Remelting of the ancient mantle domains in the Neo-Tethyan heterogeneous asthenosphere at ~130–120 Ma (context of the melting regime is modified after [Stracke, 2021](#)); (b) Subduction jump and re-initiation along detachment faults of the Neo-Tethyan ultraslow-spreading ridge and subsequent overprinting of arc-like magmatism onto the previous MORB-like oceanic crusts (modified after [Liu et al. \(2022b\)](#), [Ou et al. \(2025\)](#), and [Zhang et al. \(2019\)](#)). See the text for details.

Supplementary Table S1

[DOWNLOAD](#)

Pre-proof